

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 September 2003 (25.09.2003)

PCT

(10) International Publication Number
WO 03/078626 A2

(51) International Patent Classification⁷: **C12N 15/10**

(21) International Application Number: **PCT/DK03/00174**

(22) International Filing Date: **14 March 2003 (14.03.2003)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
PA 2002 0415 15 March 2002 (15.03.2002) DK
60/364,056 15 March 2002 (15.03.2002) US
02/00419 20 June 2002 (20.06.2002) DK
10/175,539 20 June 2002 (20.06.2002) US
60/434,427 19 December 2002 (19.12.2002) US

[DK/DK]; Slotsfogedvej 3, st. th., DK-2400 København NV (DK). **LUNDORF, Mikkel, Dybro** [DK/DK]; Charlotte Munksvej 31, 2. tv., DK-2400 København NV (DK). **HUSEMOEN, Gitte, Nystrup** [DK/DK]; Bregnerødgade 18, 1.th., DK-2200 København N (DK). **FRANCH, Thomas** [DK/DK]; Humlebækgade 14, st. tv., DK-2200 København N (DK). **THISTED, Thomas** [DK/DK]; Fjordskrænten 14, DK-3600 Frederikssund (DK).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **NUEVO-LUTION A/S** [DK/DK]; Rønnegade 8, 5th floor, DK-2100 Copenhagen Ø (DK).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GOULIAEV, Alex, Haahr** [DK/DK]; Brøndsted 223, DK-3670 Veksø Sjælland (DK). **HO, Justin** [US/DK]; Mattæusgade 50, 3.-26, DK-1666 København V (DK). **FELDING, Jakob** [DK/DK]; Ordruphøjvej 24, 1., DK-2920 Charlottenlund (DK). **SAMS, Christian** [DK/DK]; Jakob Dannefærdsvej 4A, 1., DK-1973 Frederiksberg C (DK). **PEDERSEN, Henrik** [DK/DK]; Frødesvej 24, DK-2880 Bagsværd (DK). **JENSEN, Kim, Birkebæk** [DK/DK]; Voldumvej 30C, DK-2610 Rødovre (DK). **HANSEN, Anders, Holm**

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **A BUILDING BLOCK CAPABLE OF TRANSFERRING A FUNCTIONAL ENTITY**

(57) Abstract: A building block having the dual capabilities of transferring the genetic information *e.g.* by recognising an encoding element and transferring a functional entity to a recipient reactive group is disclosed. The building block can be designed with an adjustable transferability taking into account the components of the building block. The building block may be used in the generation of a single complex or libraries of different complexes, wherein the complex comprises an encoded molecule linked to an encoding element. Libraries of complexes are useful in the quest for pharmaceutically active compounds.

WO 03/078626 A2

Title

A BUILDING BLOCK CAPABLE OF TRANSFERRING A FUNCTIONAL ENTITY

Technical Field of the Invention

5 The present invention relates to a building block comprising a complementing element and precursor for a functional entity. The building block is designed to transfer the functional entity with an adjustable efficiency to a recipient reactive group upon recognition between the complementing element and an encoding element associated with the reactive group. The invention also relates to a linkage between the
10 functional entity and the complementing element as well as a method for transferring a functional entity to recipient reactive group.

Background

15 The transfer of a chemical entity from one mono-, di- or oligonucleotide to another has been considered in the prior art. Thus, N. M. Chung *et al.* (Biochim. Biophys. Acta, 1971, 228, 536-543) used a poly(U) template to catalyse the transfer of an acetyl group from 3'-O-acetyladenosine to the 5'-OH of adenosine. The reverse transfer, i.e. the transfer of the acetyl group from a 5'-O-acetyladenosine to a 3'-OH group of another adenosine, was also demonstrated.

20 Walder *et al.* Proc. Natl. Acad. Sci. USA, 1979, 76, 51-55 suggest a synthetic procedure for peptide synthesis. The synthesis involves the transfer of nascent immobilized polypeptide attached to an oligonucleotide strand to a precursor amino acid attached to an oligonucleotide. The transfer comprises the chemical attack of the
25 amino group of the amino acid precursor on the substitution labile peptidyl ester, which in turn results in an acyl transfer. It is suggested to attach the amino acid precursor to the 5' end of an oligonucleotide with a thiol ester linkage.


30 The transfer of a peptide from one oligonucleotide to another using a template is disclosed in Bruick RK *et al.* Chemistry & Biology, 1996, 3:49-56. The carboxy terminal of the peptide is initially converted to a thioester group and subsequently transformed to an activated thioester upon incubation with Ellman's reagent. The activated thioester is reacted with a first oligo, which is 5'-thiol-terminated, resulting in the formation of a thio-ester linked intermediate. The first oligonucleotide and a
35 second oligonucleotide having a 3' amino group is aligned on a template such that

Summary of the Invention

15 Complementing Element – Linker – Carrier – C-F-connecting group – Functional entity precursor

Complementing Element is a group identifying the functional entity precursor,

Carrier is selected among the groups


$$W = \text{CH or N}$$

30 $R^2 = -H, -\text{Halogen}, -\text{NO}_2, -\text{CN}, -\text{C}(\text{Halogen})_3, -\text{C}(\text{O})\text{R}^3, -\text{C}(\text{O})\text{NHR}^3, \text{C}(\text{O})\text{NR}^3,$
 $-\text{NC}(\text{O})\text{R}^3, -\text{S}(\text{O})_2\text{NHR}^3, -\text{S}(\text{O})_2\text{NR}^3, -\text{S}(\text{O})_2\text{R}^3, -\text{P}(\text{O})_2\text{R}^3, -\text{P}(\text{O})-\text{R}^3, -\text{S}(\text{O})-\text{R}^3,$
 $\text{P}(\text{O})-\text{OR}^3, -\text{S}(\text{O})-\text{OR}^3, -\text{N}^+\text{R}^3,$ wherein p is an integer of 0 to 3, $\text{R}^3 = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl},$
 $\text{C}_1\text{-C}_6 \text{ alkenyl}, \text{C}_1\text{-C}_6 \text{ alkynyl}, \text{ or aryl}, \text{ and Halogen is F, Cl, Br, or I,}$

Y = absent, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, or -SO₂CH₂-,

C-F-connecting group is $\text{---Z---}\overset{\text{V}}{\underset{\text{X}}{\text{C}}}\text{---}$ or $\text{---}\overset{\text{V}}{\underset{\text{X}}{\text{C}}}\text{---}$ where the carrier is connected to the left hand side of the formulae and

5 X = -C-, -S-, -P-, -S(O)- or -P(O)-,

V = O, S, NH, or N-C₁-C₆ alkyl, and

Z = O, S, and

Functional entity precursor is H or selected among the group consisting of a C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₆ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, and heteroaryl, said group being substituted with 0-3 R⁴, 0-3 R⁵ and 0-3 R⁹, or selected among the group consisting of C₁-C₃ alkylene-NR⁴₂, C₁-C₃ alkylene-NR⁴C(O)R⁸, C₁-C₃ alkylene-NR⁴C(O)OR⁸, C₁-C₂ alkylene-O-NR⁴₂, C₁-C₂ alkylene-O-NR⁴C(O)R⁸, and C₁-C₂ alkylene-O-NR⁴C(O)OR⁸ substituted with 0-3 R⁹.

15 where R⁴ is H or selected independently among the group consisting of C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R⁹ and

R⁵ is selected independently from -N₃, -CNO, -C(NOH)NH₂, -NHOH, -NHNHR⁶, -C(O)R⁶, -SnR⁶₃, -B(OR⁶)₂, -P(O)(OR⁶)₂ or the group consisting of C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₆ alkadienyl said group being substituted with 0-2 R⁷,

20 where R⁶ is selected independently from H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, aryl or C₁-C₆ alkylene-aryl substituted with 0-5 halogen atoms selected from -F, -Cl, -Br, and -I; and R⁷ is independently selected from -NO₂, -COOR⁶, -COR⁶, -CN, -OSiR⁶₃, -OR⁶ and -NR⁶₂.

25 R⁸ is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₇ cycloalkyl, aryl or C₁-C₆ alkylene-aryl substituted with 0-3 substituents independently selected from -F, -Cl, -NO₂, -R³, -OR³, -SiR³₃

R⁹ is =O, -F, -Cl, -Br, -I, -CN, -NO₂, -OR⁶, -NR⁶₂, -NR⁶-C(O)R⁸, -NR⁶-C(O)OR⁸, -SR⁶, -S(O)R⁶, -S(O)₂R⁶, -COOR⁶, -C(O)NR⁶₂ and -S(O)₂NR⁶₂.

30 In the present description and claims, the direction of connections between the various components of a building block should be read left to right. For example an S-C-connecting group -C(=O)-NH- is connected to a Spacer through the carbon atom on the left and to a Carrier through the nitrogen atom on the right hand side.

The term "C₃-C₇ cycloheteroalkyl" as used herein refers to a radical of totally saturated heterocycle like a cyclic hydrocarbon containing one or more heteroatoms selected from nitrogen, oxygen, phosphor, boron and sulphur independently in the cycle such as pyrrolidine (1- pyrrolidine; 2- pyrrolidine; 3- pyrrolidine; 4- pyrrolidine; 5- pyrrolidine); pyrazolidine (1- pyrazolidine; 2- pyrazolidine; 3- pyrazolidine; 4- pyrazolidine; 5- pyrazolidine); imidazolidine (1- imidazolidine; 2- imidazolidine; 3- imidazolidine; 4- imidazolidine; 5- imidazolidine); thiazolidine (2- thiazolidine; 3- thiazolidine; 4- thiazolidine; 5- thiazolidine); piperidine (1- piperidine; 2- piperidine; 3- piperidine; 4- piperidine; 5- piperidine; 6- piperidine); piperazine (1- piperazine; 2- piperazine; 3- piperazine; 4- piperazine; 5- piperazine; 6- piperazine); morpholine (2- morpholine; 3- morpholine; 4- morpholine; 5- morpholine; 6- morpholine); thiomorpholine (2- thiomorpholine; 3- thiomorpholine; 4- thiomorpholine; 5- thiomorpholine; 6- thiomorpholine); 1,2-oxathiolane (3-(1,2-oxathiolane); 4-(1,2-oxathiolane); 5-(1,2-oxathiolane); 1,3-dioxolane (2-(1,3-dioxolane); 4-(1,3-dioxolane); 5-(1,3-dioxolane); tetrahydropyrane; (2-tetrahydropyrane; 3-tetrahydropyrane; 4-tetrahydropyrane; 5-tetrahydropyrane; 6-tetrahydropyrane); hexahydropyridazine (1-(hexahydropyridazine); 2-(hexahydropyridazine); 3-(hexahydropyridazine); 4-(hexahydropyridazine); 5-(hexahydropyridazine); 6-(hexahydropyridazine)), [1,3,2]dioxaborolane, [1,3,6,2]dioxazaborocane

The term "aryl" as used herein includes carbocyclic aromatic ring systems of 5-7 carbon atoms. Aryl is also intended to include the partially hydrogenated derivatives of the carbocyclic systems as well as up to four fused aromatic- or partially hydrogenated rings, each ring comprising 5-7 carbon atoms.

The term "heteroaryl" as used herein includes heterocyclic unsaturated ring systems containing, in addition to 2-18 carbon atoms, one or more heteroatoms selected from nitrogen, oxygen and sulphur such as furyl, thienyl, pyrrolyl, heteroaryl is also intended to include the partially hydrogenated derivatives of the heterocyclic systems enumerated below.

The terms "aryl" and "heteroaryl" as used herein refers to an aryl which can be optionally substituted or a heteroaryl which can be optionally substituted and includes phenyl, biphenyl, indenyl, naphthyl (1-naphthyl, 2-naphthyl), N-hydroxytetrazolyl, N-hydroxytriazolyl, N-hydroxyimidazolyl, anthracenyl (1-anthracenyl, 2-anthracenyl, 3-anthracenyl), thiophenyl (2-thienyl, 3-thienyl), furyl (2-furyl, 3-furyl), indolyl, oxadiazolyl, isoxazolyl, quinazolinyl, fluorenyl, xanthenyl,

isoindanyl, benzhydryl, acridinyl, thiazolyl, pyrrolyl (2-pyrrolyl), pyrazolyl (3-pyrazolyl), imidazolyl (1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl), triazolyl (1,2,3-triazol-1-yl, 1,2,3-triazol-2-yl, 1,2,3-triazol-4-yl, 1,2,4-triazol-3-yl), oxazolyl (2-oxazolyl, 4-oxazolyl, 5-oxazolyl), thiazolyl (2-thiazolyl, 4-thiazolyl, 5-thiazolyl), pyridyl (2-pyridyl, 3-pyridyl, 4-pyridyl), pyrimidinyl (2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl), pyrazinyl, pyridazinyl (3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl), quinolyl (2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl), isoquinolyl (1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl),
 benzo[b]furanyl (2-benzo[b]furanyl, 3-benzo[b]furanyl, 4-benzo[b]furanyl, 5-benzo[b]furanyl, 6-benzo[b]furanyl, 7-benzo[b]furanyl), 2,3-dihydro-benzo[b]furanyl (2-(2,3-dihydro-benzo[b]furanyl), 3-(2,3-dihydro-benzo[b]furanyl), 4-(2,3-dihydro-benzo[b]furanyl), 5-(2,3-dihydro-benzo[b]furanyl), 6-(2,3-dihydro-benzo[b]furanyl), 7-(2,3-dihydro-benzo[b]furanyl), benzo[b]thiophenyl (2-benzo[b]thiophenyl, 3-benzo[b]thiophenyl, 4-benzo[b]thiophenyl, 5-benzo[b]thiophenyl, 6-benzo[b]thiophenyl, 7-benzo[b]thiophenyl), 2,3-dihydro-benzo[b]thiophenyl (2-(2,3-dihydro-benzo[b]thiophenyl), 3-(2,3-dihydro-benzo[b]thiophenyl), 4-(2,3-dihydro-benzo[b]thiophenyl), 5-(2,3-dihydro-benzo[b]thiophenyl), 6-(2,3-dihydro-benzo[b]thiophenyl), 7-(2,3-dihydro-benzo[b]thiophenyl), indolyl (1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl), indazole (1-indazolyl, 3-indazolyl, 4-indazolyl, 5-indazolyl, 6-indazolyl, 7-indazolyl), benzimidazolyl (1-benzimidazolyl, 2-benzimidazolyl, 4-benzimidazolyl, 5-benzimidazolyl, 6-benzimidazolyl, 7-benzimidazolyl, 8-benzimidazolyl), benzoxazolyl (1-benzoxazolyl, 2-benzoxazolyl), benzothiazolyl (1-benzothiazolyl, 2-benzothiazolyl, 4-benzothiazolyl, 5-benzothiazolyl, 6-benzothiazolyl, 7-benzothiazolyl), carbazolyl (1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl), 5H-dibenz[b,f]azepine (5H-dibenz[b,f]azepin-1-yl, 5H-dibenz[b,f]azepine-2-yl, 5H-dibenz[b,f]azepine-3-yl, 5H-dibenz[b,f]azepine-4-yl, 5H-dibenz[b,f]azepine-5-yl), 10,11-dihydro-5H-dibenz[b,f]azepine (10,11-dihydro-5H-dibenz[b,f]azepine-1-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-2-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-3-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-4-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-5-yl).


The Functional Entity carries elements used to interact with host molecules and optionally reactive elements allowing further elaboration of an encoded molecule of a library. Interaction with host molecules like enzymes, receptors and polymers is typi-

cally mediated through van der waal's interactions, polar- and ionic interactions and pi-stacking effects. Substituents mediating said effects may be masked by methods known to an individual skilled in the art (Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*; 3rd ed.; John Wiley & Sons: New York, 1999.) to avoid undesired interactions or reactions during the preparation of the individual building blocks and during library synthesis. Analogously, reactive elements may be masked by suitably selected protection groups. It is appreciated by one skilled in the art that by suitable protection, a functional entity may carry a wide range of substituents.

The Functional Entity Precursor may be masked Functional Entity that is incorporated into an encoded molecule. After incorporation, reactive elements of the Functional Entity may be revealed by un-masking allowing further synthetic operations. Finally, elements mediating recognition of host molecules may be un-masked.

The function of the carrier is to adjust the transferability of the functional entity, playing the role of a leaving group. Substituents on the carrier alter the leaving group efficiency. The stronger the electron withdrawing effect the easier the functional entity is cleaved from the remainder of the building block. However the cleavage can occur too fast which will result in unspecific transfer or hydrolysis. To adjust the transferability a skilled chemist can design suitable substitutions of the carrier by evaluation of initial attempts. The transferability may be adjusted in response to the chemical composition of the functional entity, to the nature of the complementing element, to the conditions under which the transfer and recognition is performed, ect.

According to a preferred embodiment of the invention the carrier is of the general formula:




wherein W, Y, R², and p are as defined above. The transferability of the functional entity can be adjusted by suitable selection of the ring member. When the identity of W are fixed the transferability of the carrier may be adjusted by selecting type, position and amount of the ring substituents R². As an example, an unsubstituted ben-

zene ring ($W = \text{CH}$ for the entire ring structure) may be provided with an increased ability to transfer a functional entity by attaching a *Cl* in the *ortho* position. The ability to transfer functional entities may also be adjusted by proper selection of one, two or three nitrogen atoms in the ring structure. Finally, the identity and position of *Y* or alternatively the S-C-connecting group may have an influence of the transferability of the functional entity. Thus, attaching a carbonyl at the *para* position of the ring structure relative to the attachment point of the functional C-F-connecting group confers an increased ability to transfer the functional entity over a position in e.g. the *meta* position.

10

In a preferred aspect of the invention the carrier is



and attaches to the linker through *Y* and

15

$W = \text{CH}$


$R^2 = -\text{H}$, halogen, $-\text{NO}_2$, $-\text{CN}$, $-\text{C}(\text{Halogen})_3$, $-\text{C}(\text{O})\text{R}^3$, $-\text{C}(\text{O})\text{NHR}^3$, $\text{C}(\text{O})\text{NR}^3_2$, $-\text{S}(\text{O})_2\text{NHR}^3$, $-\text{S}(\text{O})_2\text{NR}^3_2$, $-\text{S}(\text{O})_2\text{R}^3$, $-\text{N}^+\text{R}^3_3$, wherein halogen is selected from the group consisting of $-\text{Cl}$, $-\text{F}$, $-\text{Br}$, and $-\text{I}$, *p* is an integer of 0 to 3, and $\text{R}^3 = \text{H}$, $\text{C}_1\text{-C}_6$ alkyl, or aryl,

20

Y = absent, $\text{C}_1\text{-C}_6$ Alkylene, or carbonyl.


The spacer serves to distance the functional entity to be transferred from the bulky complementing element. Thus, when present, the identity of the spacer is not crucial for the function of the building block. It may be desired to have a spacer which can be cleaved by light. In this occasion, the spacer is provided with e.g. the group

25




30


In the event an increased hydrophilicity is desired the spacer may be provided with a polyethylene glycol part of the general formula:



In a certain aspect of the invention the **Spacer** is a valence bond, C₁-C₈ alkylene-A-, C₁-C₈ alkenylene-A-, C₂-C₆ alkynylene-A-, or




said spacer optionally being connected through A to a linker selected from




10 where A is $\text{---C(O)NR}^1\text{---}$, $\text{---NR}^1\text{---}$, ---O--- , ---S--- , or ---C(O)---O--- ; B is ---O--- , ---S--- , $\text{---NR}^1\text{---}$ or $\text{---C(O)NR}^1\text{---}$ and connects to S-C-connecting group; R¹ is selected independently from H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₁-C₆ alkylene-aryl, or aryl substituted with 0-5 halogen atoms selected from -F, -Cl, -Br and -I; and n and m independently are integers ranging from 1 to 10.

15 More preferred the **Spacer** is C₁-C₈ alkylene-A-, C₁-C₈ alkenylene-A-, C₂-C₆ alkynylene-A-, or




said spacer optionally being connected through A to a moiety selected from




where A is $\text{---C(O)NR}^1\text{---}$, or ---S--- ; B is ---S--- , $\text{---NR}^1\text{---}$ or $\text{---C(O)NR}^1\text{---}$ and connects to S-C-connecting group; R¹ is selected independently from H, C₁-C₆ alkyl, C₁-C₆ alkylene-aryl, or aryl; and n and m independently are integers ranging from 1 to 6.

25 In certain other aspects of the invention the **Spacer** is -A-, a group C₁-C₈ alkylene-A-, C₂-C₆ alkenylene-A-, or C₂-C₆ alkynylene-A- optionally substituted with 1 to 3 hydroxy groups, or




said spacer being connected through A to a linker selected from



- 5 where A is a valence bond, $-\text{NR}^{10}$, $-\text{C}(\text{O})\text{NR}^{10}$, $-\text{NR}^{10}-\text{C}(\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{C}(\text{O})-\text{O}-$ or $-\text{OP}(=\text{O})(\text{O}^-)-\text{O}-$; B is a valence bond, $-\text{O}-$, $-\text{S}-$, $-\text{NR}^{10}$, $-\text{C}(\text{O})-$ or $-\text{C}(\text{O})\text{NR}^{10}$ - and connects to S-C-connecting group; R^{10} is selected independently from H, C_1 - C_8 al-

10 kyl, C_3 - C_7 cycloalkyl, aryl, C_1 - C_6 alkylene-aryl, $(\text{CH}_2\text{CH}_2\text{O})_n$ or $(\text{CH}_2\text{CH}_2\text{N}^+\text{G})_n$; G is H or C_1 - C_6 alkyl; and n and m independently are integers ranging from 1 to 10.

In a preferred aspect of the invention, the spacer is C_2 - C_8 alkenylene-A, said spacer being connected through A to a moiety selected from




- 15 where A is a valence bond, $-\text{C}(\text{O})\text{NR}^{10}$, $-\text{NR}^{10}-\text{C}(\text{O})-$, $-\text{S}-$, $-\text{C}(\text{O})-\text{O}-$ or $-\text{OP}(=\text{O})(\text{O}^-)-\text{O}-$; B is a valence bond, $-\text{S}-$, $-\text{NR}^{10}$, or $-\text{C}(\text{O})-$ and connects to S-C-connecting group; n and m independently are integers ranging from 1 to 10 and


20 R^{10} is selected independently from H, $(\text{CH}_2\text{CH}_2\text{O})_n$ or $(\text{CH}_2\text{CH}_2\text{N}^+\text{G})_n$, wherein G is H or C_1 - C_6 alkyl; and the spacer is connected to the complementing element through a nucleobase.

Usually, the spacer connects to the 5 position of a pyrimidine or the 7 position of a purine or deaza-purine. However, other attachment point on the nucleobase may be contemplated.

- 25 In another preferred aspect the spacer connects to the back bone of the complementing element. In this case the spacer is $-\text{A}-$,



said spacer being connected through A to a moiety selected from



where A is a valence bond, $-NR^{10}-C(O)-$, $-O-$, or $-S-$; B is a valence bond, $-S-$, $-NR^{10}-$, or $-C(O)-$ and connects to S-C-connecting group;

n and m independently are integers ranging from 1 to 10 and

R^{10} is selected independently from H, or , wherein G is H or C_1-C_6 alkyl; and the spacer is connected to the complementing element via a phosphorus group.

The phosphorus group is preferably a phosphate or a thiophosphate group attached to a 3' or a 5' end of a complementing element.



In a preferred embodiment, the complementing element serves the function of transferring genetic information *e.g.* by recognising a coding element. The recognition implies that the two parts are capable of interacting in order to assemble a complementing element – coding element complex. In the biotechnological field a variety of interacting molecular parts are known which can be used according to the invention. Examples include, but are not restricted to protein-protein interactions, protein-polysaccharide interactions, RNA-protein interactions, DNA-DNA interactions, DNA-RNA interactions, RNA-RNA interactions, biotin-streptavidin interactions, enzyme-ligand interactions, antibody-ligand interaction, protein-ligand interaction, *ect.*

The interaction between the complementing element and coding element may result in a strong or a weak bonding. If a covalent bond is formed between the parties of the affinity pair the binding between the parts can be regarded as strong, whereas the establishment of hydrogen bondings, interactions between hydrophobic domains, and metal chelation in general results in weaker bonding. In general relatively weak bonding is preferred. In a preferred aspect of the invention, the complementing element is capable of reversible interacting with the coding element so as to

provide for an attachment or detachment of the parts in accordance with the changing conditions of the media.

- In a preferred aspect of the invention, the interaction is based on nucleotides, i.e. the complementing element is a nucleic acid. Preferably, the complementing element is a sequence of nucleotides and the coding element is a sequence of nucleotides capable of hybridising to the complementing element. The sequence of nucleotides carries a series of nucleobases on a backbone. The nucleobases may be any chemical entity able to be specifically recognized by a complementing entity. The nucleobases are usually selected from the natural nucleobases (adenine, guanine, uracil, thymine, and cytosine) but also the other nucleobases obeying the Watson-Crick hydrogen-bonding rules may be used, such as the synthetic nucleobases disclosed in US 6,037,120. Examples of natural and non-natural nucleobases able to perform a specific pairing are shown in Figure 2. The backbone of the sequence of nucleotides may be any backbone able to aggregate the nucleobases is a sequence. Examples of backbones are shown in figure 4. In some aspects of the invention the addition of non-specific nucleobases to the complementing element is advantageous, figure 3.
- The coding element can be an oligonucleotide having nucleobases which complement and is specifically recognised by the complementing element, i.e. in the event the complementing element contains cytosine, the coding element part contains guanine and visa versa, and in the event the complementing element contains thymine or uracil the coding element contains adenine.
- The complementing element may be a single nucleobase. In the generation of a library, this will allow for the incorporation of four different functional entities into the template-directed molecule. However, to obtain a higher diversity a complementing element preferably comprises at least two and more preferred at least three nucleotides. Theoretically, this will provide for 4^2 and 4^3 , respectively, different functional entities uniquely identified by the complementing element. The complementing element will usually not comprise more than 100 nucleotides. It is preferred to have complementing elements with a sequence of 3 to 30 nucleotides.
- The spacer part of the linker is attached to the carrier through a S-C-connecting group (short for Spacer-Carrier-connecting group). The S-C-connecting may have any chemical composition which provides for an attachment of the Spacer with the

carrier. In certain aspect of the invention the **S-C-connecting group** is a valence bond, $-\text{NH}-\text{C}(=\text{O})-$, $-\text{NH}-\text{C}(=\text{O})-\text{C}_1-\text{C}_6$ alkylene-, $-\text{S}-\text{S}-$, $-\text{S}-\text{S}-\text{C}_1-\text{C}_6$ alkylene-, $-\text{C}_1-\text{C}_6$



5

$-\text{NH}-\text{C}(=\text{O})-\text{Arylene}-\text{C}(\text{R}^{10})_2-\text{NH}-\text{C}(=\text{O})-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{O})-\text{C}_1-\text{C}_6$ alkylene- or $-\text{C}(=\text{O})-\text{Arylene}-\text{C}(\text{R}^{10})_2-\text{NR}^{10}-\text{C}(=\text{O})-$, where the right hand side of the formulae connects to the carrier.

10 In a preferred aspect the **S-C-connecting group** is $-\text{S}-\text{S}-$, $-\text{C}_1-\text{C}_6$ alkylene- $\text{S}-\text{S}-$, $-\text{C}(=\text{O})-\text{NH}-(\text{C}_1-\text{C}_6 \text{ alkylene})-$, $-\text{C}(=\text{O})-$, or $-\text{C}(=\text{O})-\text{Arylene}-\text{C}(\text{R}^{10})_2-\text{NR}^{10}-\text{C}(=\text{O})-$, where the right hand side of the formulae connects to the carrier.

In a still more preferred aspect the **S-C-connecting group** is a valence bond, $-\text{NH}-\text{C}(=\text{O})-$, $-\text{S}-\text{S}-$, or $-\text{C}(=\text{O})-\text{NH}-$, where the right hand side of the formulae connects to the carrier.

15

The building blocks of the present invention can be used in a method for transferring a functional entity to a recipient reactive group, said method comprising the steps of providing one or more building blocks as described above and

20 contacting the one or more building blocks with a corresponding encoding element associated with a recipient reactive group under conditions which allow for a recognition between the one or more complementing elements and the encoding elements, said contacting being performed prior to, simultaneously with, or subsequent to a transfer of the functional entity to the recipient reactive group.

25

The encoding element may comprise one, two, three or more codons, i.e. sequences that may be specifically recognised by a complementing element. Each of

the codons may be separated by a suitable spacer group. Preferably, all or at least a majority of the codons of the template are arranged in sequence and each of the codons are separated from a neighbouring codon by a spacer group. Generally, it is preferred to have more than two codons on the template to allow for the synthesis of more complex encoded molecules. In a preferred aspect of the invention the number of codons of the encoding element is 2 to 100. Still more preferred are encoding elements comprising 3 to 10 codons. In another aspect, a codon comprises 1 to 50 nucleotides and the complementing element comprises a sequence of nucleotides complementary to one or more of the encoding sequences.

The recipient reactive group may be associated with the encoding element in any appropriate way. Thus, the reactive group may be associated covalently or non-covalently to the encoding element. In one embodiment the recipient reactive group is linked covalently to the encoding element through a suitable linker which may be separately cleavable to release the reaction product. In another embodiment, the reactive group is coupled to a complementing element, which is capable of recognising a sequence of nucleotides on the encoding element, whereby the recipient reactive group becomes attached to the encoding element by hybridisation. Also, the recipient reactive group may be part of a chemical scaffold, i.e. a chemical entity having one or more reactive groups available for receiving a functional entity from a building block.

The recipient reactive group may be any group able to cleave the C-F-connecting group to release the functional entity. Usually, the reactive group is nucleophilic, such as a hydroxyl, a thiol, an amine ect. A preferred recipient reactive group is an amine group. The nucleophile usually attacks the C-F-connecting group between Z and X=V or between the carrier and X=V, thereby causing the carrier group with an optional Z group to be the leaving group of the reaction and transferring the X(=V)-Functional entity precursor to the recipient. The chemical structure formed has, in the event the nucleophilic group is an amine attached to a scaffold, the general formula:

Scaffold-NH-X(=V)-Functional entity precursor

In which


X = -C-, -S-, -P-, -S(O)-, -P(O)-, and
V = O, S, NH, N-C₁-C₆ alkyl.

In a preferred aspect X is -C- and V is O.



5

The conditions which allow for transfer to occur are dependent upon the building block, notable the carrier and the C-F-connecting group, as well as the receiving reactive group. Below various examples of the conditions for a transfer to occur are depicted together with the reaction product formed.


10

A. Acylating building blocks - principle


5

B. Amide formation by reaction of amines with activated esters**C. Pyrazolone formation by reaction of hydrazines with β -Ketoesters**


10

D. Isoxazolone formation by reaction of hydroxylamines with β -Ketoesters

15


E. Pyrimidine formation by reaction of thioureas with β -Ketoesters

F. Pyrimidine formation by reaction of ureas with Malonates




5

G. Coumarine or quinolinon formation by a Heck reaction followed by a nucleophilic substitution




10


H. Phthalhydrazide formation by reaction of Hydrazines and Phthalimides



15

I. Diketopiperazine formation by reaction of Amino Acid Esters



J. Hydantoin formation by reaction of Urea and α -substituted Esters

5

According to a preferred aspect of the invention the building blocks are used for the formation of a library of compounds. The complementing element of the building block is used to identify the functional entity. Due to the enhanced proximity between reactive groups when the complementing entity and the encoding element are contacted, the functional entity together with the identity programmed in the complementing element is transferred to the encoding element associated with recipient reactive group. Thus, it is preferred that the sequence of the complementing element is unique in the sense that the same sequence is not used for another functional entity. The unique identification of the functional entity enable the possibility of decoding the encoding element in order to determine the synthetic history of the molecule formed. In the event two or more functional entities have been transferred to a scaffold, not only the identity of the transferred functional entities can be determined. Also the sequence of reaction and the type of reaction involved can be determined by decoding the encoding element. Thus, according to a preferred embodiment of the invention, each different member of a library comprises a complementing element having a unique sequence of nucleotides, which identifies the functional entity.

Brief description of the drawings

Fig. 1 shows to setups for functional entity transfer.

Fig. 2 shows examples of specific base pairing.

Fig. 3 shows examples of non-specific base-pairing

Fig. 4 shows examples of backbones.

Fig. 5 shows a gel with the results of the experiments reported in example 22.

Fig. 6 shows three examples of building block according to the present invention.

Detailed Description of the Invention

A building block of the present invention is characterized by its ability to transfer its functional entity to a receiving chemical entity. This is done by forming a new covalent bond between the receiving chemical entity and cleaving the bond between the carrier moiety and the functional entity of the building block.

Two setups for generalized functional entity transfer from a building block are depicted in figure 1. In the first example, one complementing element of a building block recognizes a template carrying another functional entity, hence bringing the functional entities in close proximity. This results in a reaction between functional entity precursor 1 and 2 forming a covalent bond between these concurrent with the cleavage of the bond between functional entity precursor 2 and its linker. In the second example, a template brings together two building blocks resulting in functional entity transfer from one building block to the other.

Fig. 6 discloses three examples of building blocks. For illustrative purposes the individual features used in the claims are indicated. In the upper compound the spacer part of the linker connects to a 3'-phosphate group of an oligonucleotide. The first part of the linker, i.e. the spacer, is an aliphatic chain ending in a nitrogen atom. The nitrogen atom bridges to the S-C-connecting group, which is an N-acylated arylmethyleamine. The carrier attached to the left hand side carbonyl group of the S-C-connecting group is a nitrophenyl group. In the para position of the nitrophenyl group, the C-F-connecting group is attached. When the building block is presented to a nucleophilic group, the functional entity precursor and the carbonyl group of the C-F-connecting group is transferred. In the event the nucleophilic group is an amine, the bond formed is an amide bond.

The middle compound of Fig. 6 discloses a linker attached to the 5' position of an oligonucleotide. The linker is attached through a 5' phosphate group and extends into a short 3 member aliphatic chain to another phosphate group which is connected to a linker terminal nitrogen group via a PEG part. The linker nitrogen group is connected to the carrier via a carbonyl group. The carrier is of the thiophenyl type as the sulphur of the C-F-connecting group connects to the ring structure. When the building block is presented to a nucleophilic group, such as an amine, the functional entity precursor together with the carbonyl group of the C-F-connecting group is

transferred to said recipient group forming an amide bond when the nucleophile is an amine.

5 The lower compound shown on Fig. 6 illustrates an example of the linker being connected to the nucleobase of the oligonucleotide complementing element. More specifically, the linker connects to the 5 position of a pyrimidine. The linker extends through an $\alpha - \beta$ unsaturated N-methylated amide to the S-C-connecting group, which is a 4-amino methyl benzoic acid derivative. The carrier is of the phenol type and the functional entity precursor together with the thiocarbonyl group of the C-F-
10 connecting group may be transferred to a recipient reactive group forming an amide in the event the recipient reactive group is an amine.


In a library synthesis, several building blocks are mixed in a reaction vessel and the added templates ensure that the building blocks - consequently the functional entities - are combined in the desired manner. As several building blocks are employed
15 at the same time, the use of *in situ* generated building blocks is disfavoured for practical reasons.

Building blocks for library synthesis should possess the necessary reactivity to enable the transfer of the functional entity but should also be stable enough to endure storage and the conditions applied during library synthesis. Hence fine tuning of the reactivity for a particular building block is vital. The reactivity of a building block depends partly on the characteristics of the functional entity and the characteristics of the carrier. E.g. a highly reactive functional entity attached to a highly reactive carrier
20 would form a building block that may be susceptible to hydrolysis during the library synthesis thus preventing successful transfer of one functional entity to another. Further, if transfer of a functional entity precursor is faster than coding element - complementing element recognition unspecific reactions may result. Therefore, the present invention particularly relates to practically useful library building blocks capable of acting as acylating agents, thioacetylating agents or amidinoylating agents with a balanced reactivity. Such building blocks may be assembled by
25 several different pathways as described below.

Formation of an amide bond between a carboxylic acid of the Carrier and an amine group of a Spacer

The Carrier-Functional Entity Precursor ensemble may be bound to the Spacer by several different reactions as illustrated below.





5







10

Examples of Carrier-Functional Entity Precursor reagents:

<p>The structure shows a 'Carrier' group (a benzene ring with substituents W, W, W, Y, and R') connected to a 'Functional Entity' (X) via a bond (Z). The carrier also has a carboxylic acid group (COOH) attached to it.</p>	<p>Z = O, S X = -C-, -S-, -P-, -S(O)-, -P(O)- V = O, S, NR, R = H, C₁-C₆ alkyl W = CH or N, chosen independently R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃, R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently Y = absent, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>
---	---

	<p>Z = O, S</p> <p>X = -C-, -S-, -P-, -S(O)-, -P(O)-</p> <p>V = O, S, NR, R = H, C₁-C₈ alkyl</p> <p>Y = absent, C₁-C₈ Alkyl, C₁-C₈ Alkenyl, C₁-C₈ Alkynyl, Aryl, Heteroaryl, Carbonyl, -SO₂CH₂-</p>
	<p>Z = S</p> <p>R' = -CH₂-</p> <p>X = -C-, -S-, -P-, -S(O)-, -P(O)-</p> <p>V = O, S, NR, R = H, C₁-C₈ alkyl</p> <p>Y = nothing, C₁-C₈ Alkylene, C₁-C₈ Alkenylene, C₁-C₈ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>
	<p>W = CH or N</p> <p>X = -C-, -S-, -P-, -S(O)-, -P(O)-</p> <p>V = O, S, NR, R = H, C₁-C₈ alkyl</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃, R'' = alkyl, alkenyl, alkynyl, aryl.</p> <p>Y = nothing, C₁-C₈ Alkyl, C₁-C₈ Alkenyl, C₁-C₈ Alkynyl, Aryl, Heteroaryl, Carbonyl, -SO₂CH₂-</p>
	<p>W = CH or N</p> <p>X = -C-, -S-, -P-, -S(O)-, -P(O)-</p> <p>V = O, S, NR, R = H, C₁-C₈ alkyl</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃, R'' = H, C₁-C₈ alkyl, C₁-C₈ alkenyl, C₁-C₈ alkynyl or aryl, chosen independently</p> <p>Y = absent, C₁-C₈ Alkylene, C₁-C₈ Alkenylene, C₁-C₈ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>


 <p>Functional Entity</p>	<p>X = -C-, -S-, -P-, -S(O)-, -P(O)-</p> <p>V = O, S, NR, R = H, C₁-C₆ alkyl</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃, R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>
 <p>Functional Entity</p>	<p>W = CH or N</p> <p>X = -C-, -S-, -P-, -S(O)-, -P(O)-</p> <p>V = O, S, NR, R = H, C₁-C₆ alkyl</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃, R'' = alkyl, alkenyl, alkynyl, aryl.</p> <p>Y = nothing, C₁-C₆ Alkyl, C₁-C₆ Alkenyl, C₁-C₆ Alkynyl, Aryl, Heteroaryl, Carbonyl, -SO₂CH₂-</p>
 <p>Functional Entity</p>	<p>Z = O, S</p> <p>X = -C-, -S-, -P-, -S(O)-, -P(O)-</p> <p>V = O, S, NR, R = H, C₁-C₆ alkyl</p> <p>W = CH or N, chosen independently</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃ chosen independently</p> <p>p = 0, 1, 2, 3 or 4</p> <p>Y = absent, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>

Stepwise loading of the carrier and the functional entity

X = leaving group

Sequential loading of the carrier and the functional entity allows other types of chemistries to be used.





5


Carrier introduced via amide bond formation

10

Examples of Carrier reactants:



	<p>Z = O, S</p> <p>W = CH or N, independently chosen</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃,</p> <p>R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>
--	---


	<p>Z = O, S</p> <p>Y = nothing, C₁-C₆ Alkyl, C₁-C₆ Alkenyl, C₁-C₆ Alkynyl, Aryl, Heteroaryl, Carbonyl, -SO₂CH₂-</p>
	<p>W = CH or N</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃,</p> <p>R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>
	<p>W = CH or N</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃,</p> <p>R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>
	<p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃,</p> <p>R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>

Carrier introduced via S-S bond formation

Examples of Carrier reactants:

	<p>Z = O</p> <p>W = CH or N</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃,</p> <p>R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>
	<p>Z = O</p> <p>Y = nothing, C₁-C₆ Alkyl, C₁-C₆ Alkenyl, C₁-C₆ Alkynyl, Aryl, Heteroaryl, Carbonyl, -SO₂CH₂-</p>
	<p>W = CH or N</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃,</p> <p>R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl,</p>

	-SO ₂ CH ₂ -
	<p>W = CH or N</p> <p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃,</p> <p>R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>
	<p>R' = -H, -Halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R'', -C(O)NHR'', C(O)NR''₂, -NC(O)R'', -S(O)₂NHR'', -S(O)₂NR''₂, -S(O)₂R'', -P(O)₂-R'', -P(O)-R'', -S(O)-R'', P(O)-OR'', -S(O)-OR'', -N⁺R''₃,</p> <p>R'' = H, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl or aryl, chosen independently</p> <p>Y = nothing, C₁-C₆ Alkylene, C₁-C₆ Alkenylene, C₁-C₆ Alkynylene, Arylene, Heteroarylene, Carbonyl, -SO₂CH₂-</p>

Functional Entity introduced as a thioacid

Lg = leaving group

Examples of Carrier reactants:

	$\text{R}^1 = -\text{CH}_2-$ $\text{Y} = \text{nothing, C}_1\text{-C}_6 \text{ Alkylene, C}_1\text{-C}_6 \text{ Alkenylene, C}_1\text{-C}_6 \text{ Alkynylene, Arylene, Heteroarylene, Carbonyl, } -\text{SO}_2\text{CH}_2-$
--	---

5

As discussed above the **C-F-connecting group** may be selected from a large group of compounds of the general formula **-Z-(X=V)-** or **-(X=V)-**. In certain aspects of the invention $\text{X} = \text{C}, \text{S}, \text{P}, \text{S}(=\text{O}),$ or $\text{P}(=\text{O})$, in another preferred embodiment $\text{X} = \text{C}, \text{S},$ or $\text{S}(=\text{O})$, and in still another preferred embodiment $\text{X} = \text{C}$. In certain aspects of the invention $\text{V} = \text{O}, \text{S}, \text{NR}^{10}$ or NOR^{10} , in another preferred embodiment $\text{V} = \text{O}$ or NR^{10} , and in still another preferred embodiment $\text{V} = \text{O}$. In a certain aspect of the invention $\text{Z} = \text{O},$ or S , in another preferred embodiment, $\text{Z} = \text{O}$, and in still another preferred embodiment, $\text{Z} = \text{S}$.

15

Wherein R^{10} is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of $\text{SnR}^{11}\text{R}^{12}, \text{R}^{13}, \text{Sn}(\text{OR}^{11})\text{R}^{12}\text{R}^{13}, \text{Sn}(\text{OR}^{11})(\text{OR}^{12})\text{R}^{13}, \text{BR}^{11}\text{R}^{12}, \text{B}(\text{OR}^{11})\text{R}^{12}, \text{B}(\text{OR}^{11})(\text{OR}^{12}),$ halogen, CN, CNO, $\text{C}(\text{halogen})_3,$ $\text{OR}^{11}, \text{OC}(=\text{O})\text{R}^{11}, \text{OC}(=\text{O})\text{OR}^{11}, \text{OC}(=\text{O})\text{NR}^{11}\text{R}^{12}, \text{SR}^{11}, \text{S}(=\text{O})\text{R}^{11}, \text{S}(=\text{O})_2\text{R}^{11}, \text{S}(=\text{O})_2\text{NR}^{11}\text{R}^{12}, \text{NO}_2, \text{N}_3, \text{NR}^{11}\text{R}^{12}, \text{N}^+\text{R}^{11}\text{R}^{12}\text{R}^{13}, \text{NR}^{11}\text{OR}^{12}, \text{NR}^{11}\text{NR}^{12}\text{R}^{13}, \text{NR}^{11}\text{C}(=\text{O})\text{R}^{12}, \text{NR}^{11}\text{C}(=\text{O})\text{OR}^{12}, \text{NR}^{11}\text{C}(=\text{O})\text{NR}^{12}\text{R}^{13}, \text{NC}, \text{P}(=\text{O})(\text{OR}^{11})\text{OR}^{12}, \text{P}^+\text{R}^{11}\text{R}^{12}\text{R}^{13}, \text{C}(=\text{O})\text{R}^{11}, \text{C}(=\text{NR}^{11})\text{R}^{12}, \text{C}(=\text{NOR}^{11})\text{R}^{12}, \text{C}(=\text{NNR}^{11}\text{R}^{12}), \text{C}(=\text{O})\text{OR}^{11}, \text{C}(=\text{O})\text{NR}^{11}\text{R}^{12}, \text{C}(=\text{O})\text{NR}^{11}\text{OR}^{12}, \text{C}(=\text{O})\text{NR}^{11}\text{NR}^{12}\text{R}^{13}, \text{C}(=\text{NR}^{11})\text{NR}^{12}\text{R}^{13}, \text{C}(=\text{NOR}^{11})\text{NR}^{12}\text{R}^{13}$ or $\text{R}^{14},$ wherein,

25

R^{11} , R^{12} and R^{13} independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of halogen, CN, CNO, C(halogen)₃, OR¹⁴, OC(=O)R¹⁴, OC(=O)OR¹⁴, OC(=O)NR¹⁴R¹⁵, SR¹⁴, S(=O)R¹⁴, S(=O)₂R¹⁴,
 5 S(=O)₂NR¹⁴R¹⁵, NO₂, N₃, NR¹⁴R¹⁵, N⁺R¹⁴R¹⁵R¹⁶, NR¹¹OR¹², NR¹¹NR¹²R¹³, NR¹⁴C(=O)R¹⁵, NR¹⁴C(=O)OR¹⁵, NR¹⁴C(=O)NR¹⁵R¹⁶, NC, P(=O)(OR¹⁴)OR¹⁵, P⁺R¹¹R¹²R¹³, C(=O)R¹⁴, C(=NR¹⁴)R¹⁵, C(=NOR¹⁴)R¹⁵, C(=NNR¹⁴R¹⁵), C(=O)OR¹⁴, C(=O)NR¹⁴R¹⁵, C(=O)NR¹⁴OR¹⁵, C(=NR¹¹)NR¹²R¹³, C(=NOR¹¹)NR¹²R¹³ or C(=O)NR¹⁴NR¹⁵R¹⁶, wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or
 10 R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring, wherein,
 R^{14} , R^{15} and R^{16} independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl and wherein R^{14} and R^{15} may together form a 3-8
 15 membered heterocyclic ring or R^{14} and R^{16} may together form a 3-8 membered heterocyclic ring or R^{15} and R^{16} may together form a 3-8 membered heterocyclic ring,

in a further preferred embodiment,
 R^{10} is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl,
 20 C₃-C₇ cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of SnR¹¹R¹²,R¹³, Sn(OR¹¹)R¹²R¹³, Sn(OR¹¹)(OR¹²)R¹³, BR¹¹R¹², B(OR¹¹)R¹², B(OR¹¹)(OR¹²), halogen, CN, CNO, C(halogen)₃, OR¹¹, OC(=O)R¹¹, OC(=O)OR¹¹, OC(=O)NR¹¹R¹², SR¹¹, S(=O)R¹¹, S(=O)₂R¹¹, S(=O)₂NR¹¹R¹², NO₂, N₃, NR¹¹R¹², N⁺R¹¹R¹²R¹³, NR¹¹OR¹², NR¹¹NR¹²R¹³,
 25 NR¹¹C(=O)R¹², NR¹¹C(=O)OR¹², NR¹¹C(=O)NR¹²R¹³, NC, P(=O)(OR¹¹)OR¹², P⁺R¹¹R¹²R¹³, C(=O)R¹¹, C(=NR¹¹)R¹², C(=NOR¹¹)R¹², C(=NNR¹¹R¹²), C(=O)OR¹¹, C(=O)NR¹¹R¹², C(=O)NR¹¹OR¹², C(=O)NR¹¹NR¹²R¹³, C(=NR¹¹)NR¹²R¹³, C(=NOR¹¹)NR¹²R¹³ or R¹⁴,
 wherein,
 30 R^{11} , R^{12} , R^{13} and R^{14} independently is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

35

in another preferred embodiment,

- R^{10} is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of halogen, CN, $C(\text{halogen})_3$, OR^{11} , $OC(=O)R^{11}$, $OC(=O)OR^{11}$, $OC(=O)NR^{11}R^{12}$,
 5 SR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}OR^{12}$, $NR^{11}NR^{12}R^{13}$,
 $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $P(=O)(OR^{11})OR^{12}$, $C(=O)R^{11}$,
 $C(=NR^{11})R^{12}$, $C(=NOR^{11})R^{12}$, $C(=NNR^{11}R^{12})$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$,
 $C(=O)NR^{11}OR^{12}$, $C(=O)NR^{11}NR^{12}R^{13}$, $C(=NR^{11})NR^{12}R^{13}$, $C(=NOR^{11})NR^{12}R^{13}$ or R^{14} ,
 wherein,
 10 R^{11} , R^{12} , R^{13} and R^{14} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

- 15 in still another preferred embodiment,

- R^{10} is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $OC(=O)R^{11}$, $OC(=O)OR^{11}$, $OC(=O)NR^{11}R^{12}$, SR^{11} , $S(=O)R^{11}$,
 $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}OR^{12}$, $NR^{11}NR^{12}R^{13}$, $NR^{11}C(=O)R^{12}$,
 20 $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $P(=O)(OR^{11})OR^{12}$, $C(=O)R^{11}$, $C(=NR^{11})R^{12}$,
 $C(=NOR^{11})R^{12}$, $C(=NNR^{11}R^{12})$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$,
 $C(=O)NR^{11}NR^{12}R^{13}$, $C(=NR^{11})NR^{12}R^{13}$, $C(=NOR^{11})NR^{12}R^{13}$ or R^{14} ,
 wherein,
 R^{11} , R^{12} , R^{13} and R^{14} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

in still another preferred embodiment,

- 30 R^{10} is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$,
 $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$,
 $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
 35 wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

5

in still another preferred embodiment,

R^{10} is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl, morpholinyl, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹¹, S(=O)R¹¹, S(=O)₂R¹¹, S(=O)₂NR¹¹R¹², NO₂, NR¹¹R¹², NR¹¹C(=O)R¹², NR¹¹C(=O)OR¹², NR¹¹C(=O)NR¹²R¹³, C(=O)R¹¹, C(=O)NR¹¹R¹², C(=O)OR¹¹, C(=O)NR¹¹R¹², C(=O)NR¹¹OR¹² or R¹⁴,

10

wherein,

15

R^{11} , R^{12} , R^{13} and R^{14} independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

20

in still another preferred embodiment,

R^{10} is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹¹, S(=O)R¹¹, S(=O)₂R¹¹, S(=O)₂NR¹¹R¹², NO₂, NR¹¹R¹², NR¹¹C(=O)R¹², NR¹¹C(=O)OR¹², NR¹¹C(=O)NR¹²R¹³, C(=O)R¹¹, C(=O)NR¹¹R¹², C(=O)OR¹¹, C(=O)NR¹¹R¹², C(=O)NR¹¹OR¹² or R¹⁴,

25

wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

30

in still another preferred embodiment,

R^{10} is H, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹¹, S(=O)R¹¹, S(=O)₂R¹¹, S(=O)₂NR¹¹R¹², NO₂, NR¹¹R¹², NR¹¹C(=O)R¹²,

35

$\text{NR}^{11}\text{C}(=\text{O})\text{OR}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{NR}^{12}\text{R}^{13}$, $\text{C}(=\text{O})\text{R}^{11}$, $\text{C}(=\text{NOR}^{11})\text{R}^{12}$, $\text{C}(=\text{O})\text{OR}^{11}$,
 $\text{C}(=\text{O})\text{NR}^{11}\text{R}^{12}$, $\text{C}(=\text{O})\text{NR}^{11}\text{OR}^{12}$ or R^{14} ,

wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$ cyclo-
 5 heteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8
 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered het-
 erocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

in still another preferred embodiment,

10 R^{10} is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally
 substituted with one or more substituents selected from the group consisting of F,
 Cl, CN, CF_3 , OR^{11} , $\text{S}(=\text{O})\text{R}^{11}$, $\text{S}(=\text{O})_2\text{R}^{11}$, $\text{S}(=\text{O})_2\text{NR}^{11}\text{R}^{12}$, NO_2 , $\text{NR}^{11}\text{R}^{12}$,
 $\text{NR}^{11}\text{C}(=\text{O})\text{R}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{OR}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{NR}^{12}\text{R}^{13}$, $\text{C}(=\text{O})\text{R}^{11}$, $\text{C}(=\text{NOR}^{11})\text{R}^{12}$,
 $\text{C}(=\text{O})\text{OR}^{11}$, $\text{C}(=\text{O})\text{NR}^{11}\text{R}^{12}$, $\text{C}(=\text{O})\text{NR}^{11}\text{OR}^{12}$ or R^{14} ,

15 wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$ cyclo-
 heteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8
 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered het-
 erocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

20

in still another preferred embodiment,

R^{10} is H, phenyl or naphthyl optionally substituted with one or more substituents se-
 lected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $\text{S}(=\text{O})\text{R}^{11}$, $\text{S}(=\text{O})_2\text{R}^{11}$,
 $\text{S}(=\text{O})_2\text{NR}^{11}\text{R}^{12}$, NO_2 , $\text{NR}^{11}\text{R}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{R}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{OR}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{NR}^{12}\text{R}^{13}$,
 25 $\text{C}(=\text{O})\text{R}^{11}$, $\text{C}(=\text{NOR}^{11})\text{R}^{12}$, $\text{C}(=\text{O})\text{OR}^{11}$, $\text{C}(=\text{O})\text{NR}^{11}\text{R}^{12}$, $\text{C}(=\text{O})\text{NR}^{11}\text{OR}^{12}$ or R^{14} ,

wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$ cyclo-
 heteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8
 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered het-
 erocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

30

in still another preferred embodiment,

R^{10} is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with
 one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} ,
 35 $\text{S}(=\text{O})\text{R}^{11}$, $\text{S}(=\text{O})_2\text{R}^{11}$, $\text{S}(=\text{O})_2\text{NR}^{11}\text{R}^{12}$, NO_2 , $\text{NR}^{11}\text{R}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{R}^{12}$,

$\text{NR}^{11}\text{C}(=\text{O})\text{OR}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{NR}^{12}\text{R}^{13}$, $\text{C}(=\text{O})\text{R}^{11}$, $\text{C}(=\text{NOR}^{11})\text{R}^{12}$, $\text{C}(=\text{O})\text{OR}^{11}$,
 $\text{C}(=\text{O})\text{NR}^{11}\text{R}^{12}$, $\text{C}(=\text{O})\text{NR}^{11}\text{OR}^{12}$ or R^{14} ,

wherein,

- 5 R^{11} , R^{12} , R^{13} and R^{14} independently is H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$ cyclo-
heteroalkyl, aryl or heteroaryl and wherein R^{11} and R^{12} may together form a 3-8
membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered het-
erocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

in still another preferred embodiment,

- 10 R^{10} is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclo-
hexyl optionally substituted with one or more substituents selected from the group
consisting of F, Cl, CN, CF_3 , OR^{11} , $\text{S}(=\text{O})\text{R}^{11}$, $\text{S}(=\text{O})_2\text{R}^{11}$, $\text{S}(=\text{O})_2\text{NR}^{11}\text{R}^{12}$, NO_2 ,
 $\text{NR}^{11}\text{R}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{R}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{OR}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{NR}^{12}\text{R}^{13}$, $\text{C}(=\text{O})\text{R}^{11}$,
 $\text{C}(=\text{NOR}^{11})\text{R}^{12}$, $\text{C}(=\text{O})\text{OR}^{11}$, $\text{C}(=\text{O})\text{NR}^{11}\text{R}^{12}$, $\text{C}(=\text{O})\text{NR}^{11}\text{OR}^{12}$ or R^{14} ,
15 wherein,
 R^{11} , R^{12} , R^{13} and R^{14} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl,
cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quino-
linyl or isoquinoliny and wherein R^{11} and R^{12} may together form a 3-8 membered
heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring
20 or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

in still another preferred embodiment,

- R^{10} is H, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substi-
tuted with one or more substituents selected from the group consisting of F, Cl, CN,
25 CF_3 , OR^{11} , $\text{S}(=\text{O})\text{R}^{11}$, $\text{S}(=\text{O})_2\text{R}^{11}$, $\text{S}(=\text{O})_2\text{NR}^{11}\text{R}^{12}$, NO_2 , $\text{NR}^{11}\text{R}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{R}^{12}$,
 $\text{NR}^{11}\text{C}(=\text{O})\text{OR}^{12}$, $\text{NR}^{11}\text{C}(=\text{O})\text{NR}^{12}\text{R}^{13}$, $\text{C}(=\text{O})\text{R}^{11}$, $\text{C}(=\text{NOR}^{11})\text{R}^{12}$, $\text{C}(=\text{O})\text{OR}^{11}$,
 $\text{C}(=\text{O})\text{NR}^{11}\text{R}^{12}$, $\text{C}(=\text{O})\text{NR}^{11}\text{OR}^{12}$ or R^{14} ,
wherein,
 R^{11} , R^{12} , R^{13} and R^{14} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl,
30 cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quino-
linyl or isoquinoliny and wherein R^{11} and R^{12} may together form a 3-8 membered
heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring
or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

- 35 in still another preferred embodiment,

R^{10} is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$,
 5 $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
 wherein,
 R^{11} , R^{12} , R^{13} and R^{14} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^{11} and R^{12} may together form a 3-8 membered
 10 heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

in still another preferred embodiment,
 R^{10} is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$,
 15 $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
 wherein,
 R^{11} , R^{12} , R^{13} and R^{14} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl,
 20 cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

25 in still another preferred embodiment,
 R^{10} is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$,
 30 $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
 wherein,
 R^{11} , R^{12} , R^{13} and R^{14} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^{11} and R^{12} may together form a 3-8 membered

heterocyclic ring or R¹¹ and R¹³ may together form a 3-8 membered heterocyclic ring or R¹² and R¹³ may together form a 3-8 membered heterocyclic ring,

in still another preferred embodiment,

5 R¹⁰ is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹¹, S(=O)R¹¹, S(=O)₂R¹¹, S(=O)₂NR¹¹R¹², NO₂, NR¹¹R¹², NR¹¹C(=O)R¹², NR¹¹C(=O)OR¹², NR¹¹C(=O)NR¹²R¹³, C(=O)R¹¹, C(=O)NR¹¹R¹², C(=O)OR¹¹, C(=O)NR¹¹R¹², C(=O)NR¹¹OR¹² or R¹⁴,

10 wherein,

R¹¹, R¹², R¹³ and R¹⁴ independently is H, methyl, ethyl, propyl or butyl and wherein R¹¹ and R¹² may together form a 3-8 membered heterocyclic ring or R¹¹ and R¹³ may together form a 3-8 membered heterocyclic ring or R¹² and R¹³ may together form a 3-8 membered heterocyclic ring,

15

in still another preferred embodiment,

R¹⁰ is H, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹¹, S(=O)R¹¹, S(=O)₂R¹¹, S(=O)₂NR¹¹R¹², NO₂, NR¹¹R¹², NR¹¹C(=O)R¹²,
20 NR¹¹C(=O)OR¹², NR¹¹C(=O)NR¹²R¹³, C(=O)R¹¹, C(=O)NR¹¹R¹², C(=O)OR¹¹, C(=O)NR¹¹R¹², C(=O)NR¹¹OR¹² or R¹⁴,

wherein,

R¹¹, R¹², R¹³ and R¹⁴ independently is H, methyl, ethyl, propyl or butyl and wherein R¹¹ and R¹² may together form a 3-8 membered heterocyclic ring or R¹¹ and R¹³ may together form a 3-8 membered heterocyclic ring or R¹² and R¹³ may together form a 3-8 membered heterocyclic ring,

25

in still another preferred embodiment,

R¹⁰ is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹¹, S(=O)R¹¹, S(=O)₂R¹¹, S(=O)₂NR¹¹R¹², NO₂, NR¹¹R¹², NR¹¹C(=O)R¹², NR¹¹C(=O)OR¹², NR¹¹C(=O)NR¹²R¹³, C(=O)R¹¹, C(=O)NR¹¹R¹², C(=O)OR¹¹, C(=O)NR¹¹R¹², C(=O)NR¹¹OR¹² or R¹⁴,
30 wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, methyl, ethyl, propyl or butyl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,

5

in still another preferred embodiment,

R^{10} is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
 10 wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, methyl, ethyl, propyl or butyl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,
 15

in still another preferred embodiment,

R^{10} is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
 20 wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, methyl, ethyl, propyl or butyl and wherein R^{11} and R^{12} may together form a 3-8 membered heterocyclic ring or R^{11} and R^{13} may together form a 3-8 membered heterocyclic ring or R^{12} and R^{13} may together form a 3-8 membered heterocyclic ring,
 25

in still another preferred embodiment,

R^{10} is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
 30

35 wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

in still another preferred embodiment,

5 R^{10} is aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,

10 wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

in still another preferred embodiment,

15 R^{10} is phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,

20 wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

in still another preferred embodiment,

25 R^{10} is phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,

wherein,

30 R^{11} , R^{12} , R^{13} and R^{14} independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

in still another preferred embodiment,

35 R^{10} is thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} ,

$S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$,
 $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$,
 $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,

wherein,

- 5 R^{11} , R^{12} , R^{13} and R^{14} independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

in still another preferred embodiment,

- 10 R^{10} is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,

wherein,

- 15 R^{11} , R^{12} , R^{13} and R^{14} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

in still another preferred embodiment,

- 20 R^{10} is aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,

wherein,

- 25 R^{11} , R^{12} , R^{13} and R^{14} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

in still another preferred embodiment,

- 30 R^{10} is phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,

wherein,

R^{11} , R^{12} , R^{13} and R^{14} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

in still another preferred embodiment,

- 5 R^{10} is phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
wherein,

- 10 R^{11} , R^{12} , R^{13} and R^{14} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

in still another preferred embodiment,

- 15 R^{10} is thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{11} , $S(=O)R^{11}$, $S(=O)_2R^{11}$, $S(=O)_2NR^{11}R^{12}$, NO_2 , $NR^{11}R^{12}$, $NR^{11}C(=O)R^{12}$, $NR^{11}C(=O)OR^{12}$, $NR^{11}C(=O)NR^{12}R^{13}$, $C(=O)R^{11}$, $C(=NOR^{11})R^{12}$, $C(=O)OR^{11}$, $C(=O)NR^{11}R^{12}$, $C(=O)NR^{11}OR^{12}$ or R^{14} ,
wherein,

- 20 R^{11} , R^{12} , R^{13} and R^{14} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

in still another preferred embodiment,

R^{10} is H, C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl

25

in still another preferred embodiment,

R^{10} is H,

in still another preferred embodiment,

- 30 R^{10} is C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl or C_3 - C_7 cycloheteroalkyl,

in still another preferred embodiment,

R^{10} is methyl, ethyl, propyl or butyl

- 35 in still another preferred embodiment

R¹⁰ is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl

in still another preferred embodiment

R¹⁰ is aziridiny, pyrrolidiny, piperidiny or morpholiny

5

in still another preferred embodiment,

R¹⁰ is aryl or heteroaryl

in still another preferred embodiment,

10

R¹⁰ is phenyl or naphthyl

in still another preferred embodiment,

R¹⁰ is thienyl, furyl, pyridyl, quinoliny or isoquinolyl.

15 The **Functional entity precursor** may be selected from any transferable chemical group capable of forming a connection to the C-F-connecting group. In certain aspects of the invention the functional entity precursor is represented by the formula Z²R¹⁷

20 wherein Z is absent, O, S or NR²⁴. In certain embodiment Z is absent. In a another embodiment Z is O. In still another embodiment Z is S, and in still a further embodiment Z is NR²⁴.

25 R¹⁷ and R²⁴ independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cyclo-heteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of SnR¹⁸R¹⁹, R²⁰, Sn(OR¹⁸)R¹⁹R²⁰, Sn(OR¹⁸)(OR¹⁹)R²⁰, BR¹⁸R¹⁹, B(OR¹⁸)R¹⁹, B(OR¹⁸)(OR¹⁹), halogen, CN, CNO, C(halogen)₃, OR¹⁸, OC(=O)R¹⁸, OC(=O)OR¹⁸, OC(=O)NR¹⁸R¹⁹, SR¹⁸, S(=O)R¹⁸, S(=O)₂R¹⁸, S(=O)₂NR¹⁸R¹⁹, NO₂, N₃, NR¹⁸R¹⁹, N⁺R¹⁸R¹⁹R²⁰, NR¹⁸OR¹⁹, NR¹⁸NR¹⁹R²⁰, NR¹⁸C(=O)R¹⁹, NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, NC, P(=O)(OR¹⁸)OR¹⁹, P⁺R¹⁸R¹⁹R²⁰, C(=O)R¹⁸, C(=NR¹⁸)R¹⁹, C(=NOR¹⁸)R¹⁹, C(=NNR¹⁸R¹⁹), C(=O)OR¹⁸, C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹, C(=O)NR¹⁸NR¹⁹R²⁰, C(=NR¹⁸)NR¹⁹R²⁰, C(=NOR¹⁸)NR¹⁹R²⁰ or R²¹,
30 wherein,

R^{18} , R^{19} and R^{20} independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of halogen, CN, CNO, C(halogen)₃, OR²¹, OC(=O)R²¹, OC(=O)OR²¹, OC(=O)NR²¹R²², SR²¹, S(=O)R²¹, S(=O)₂R²¹, S(=O)₂NR²¹R²², NO₂, N₃, NR²¹R²², N⁺R²¹R²²R²³, NR¹⁸OR¹⁹, NR¹⁸NR¹⁹R²⁰, NR²¹C(=O)R²², NR²¹C(=O)OR²², NR²¹C(=O)NR²²R²³, NC, P(=O)(OR²¹)OR²², P⁺R¹⁸R¹⁹R²⁰, C(=O)R²¹, C(=NR²¹)R²², C(=NOR²¹)R²², C(=NNR²¹R²²), C(=O)OR²¹, C(=O)NR²¹R²², C(=O)NR²¹OR²², C(=NR¹⁸)NR¹⁹R²⁰, C(=NOR¹⁸)NR¹⁹R²⁰ or C(=O)NR²¹NR²²R²³, wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

wherein,

R^{21} , R^{22} and R^{23} independently is H, alkyl, alkenyl, alkynyl, alkadienyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl and wherein R^{21} and R^{22} may together form a 3-8 membered heterocyclic ring or R^{21} and R^{23} may together form a 3-8 membered heterocyclic ring or R^{22} and R^{23} may together form a 3-8 membered heterocyclic ring,

In a further embodiment,

R^{17} and R^{24} independently is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of SnR¹⁸R¹⁹R²⁰, Sn(OR¹⁸)R¹⁹R²⁰, Sn(OR¹⁸)(OR¹⁹)R²⁰, BR¹⁸R¹⁹, B(OR¹⁸)R¹⁹, B(OR¹⁸)(OR¹⁹), halogen, CN, CNO, C(halogen)₃, OR¹⁸, OC(=O)R¹⁸, OC(=O)OR¹⁸, OC(=O)NR¹⁸R¹⁹, SR¹⁸, S(=O)R¹⁸, S(=O)₂R¹⁸, S(=O)₂NR¹⁸R¹⁹, NO₂, N₃, NR¹⁸R¹⁹, N⁺R¹⁸R¹⁹R²⁰, NR¹⁸OR¹⁹, NR¹⁸NR¹⁹R²⁰, NR¹⁸C(=O)R¹⁹, NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, NC, P(=O)(OR¹⁸)OR¹⁹, P⁺R¹⁸R¹⁹R²⁰, C(=O)R¹⁸, C(=NR¹⁸)R¹⁹, C(=NOR¹⁸)R¹⁹, C(=NNR¹⁸R¹⁹), C(=O)OR¹⁸, C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹, C(=O)NR¹⁸NR¹⁹R²⁰, C(=NR¹⁸)NR¹⁹R²⁰, C(=NOR¹⁸)NR¹⁹R²⁰ or R²¹,

wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In another embodiment,

R^{17} and R^{24} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of halogen, CN, C(halogen)₃, OR^{18} , $OC(=O)R^{18}$, $OC(=O)OR^{18}$,
 5 $OC(=O)NR^{18}R^{19}$, SR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$,
 $NR^{18}OR^{19}$, $NR^{18}NR^{19}R^{20}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$,
 $P(=O)(OR^{18})OR^{19}$, $C(=O)R^{18}$, $C(=NR^{18})R^{19}$, $C(=NOR^{18})R^{19}$, $C(=NNR^{18}R^{19})$,
 $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$, $C(=O)NR^{18}NR^{19}R^{20}$, $C(=NR^{18})NR^{19}R^{20}$,
 $C(=NOR^{18})NR^{19}R^{20}$ or R^{21} ,

10 wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

15

In still another embodiment,

R^{17} and R^{24} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $OC(=O)R^{18}$, $OC(=O)OR^{18}$,
 20 $OC(=O)NR^{18}R^{19}$, SR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$,
 $NR^{18}OR^{19}$, $NR^{18}NR^{19}R^{20}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$,
 $P(=O)(OR^{18})OR^{19}$, $C(=O)R^{18}$, $C(=NR^{18})R^{19}$, $C(=NOR^{18})R^{19}$, $C(=NNR^{18}R^{19})$,
 $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$, $C(=O)NR^{18}NR^{19}R^{20}$, $C(=NR^{18})NR^{19}R^{20}$,
 $C(=NOR^{18})NR^{19}R^{20}$ or R^{21} ,

25 wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

30

In still another embodiment,

R^{17} and R^{24} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$,

NO_2 , $\text{NR}^{18}\text{R}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{R}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{OR}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{NR}^{19}\text{R}^{20}$, $\text{C}(=\text{O})\text{R}^{18}$, $\text{C}(=\text{NOR}^{18})\text{R}^{19}$, $\text{C}(=\text{O})\text{OR}^{18}$, $\text{C}(=\text{O})\text{NR}^{18}\text{R}^{19}$, $\text{C}(=\text{O})\text{NR}^{18}\text{OR}^{19}$ or R^{21} ,

wherein,

- 5 R^{18} , R^{19} , R^{20} and R^{21} independently is H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$ cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

- 10 R^{17} and R^{24} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl, morpholinyl, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $\text{S}(=\text{O})\text{R}^{18}$, $\text{S}(=\text{O})_2\text{R}^{18}$, $\text{S}(=\text{O})_2\text{NR}^{18}\text{R}^{19}$, NO_2 , $\text{NR}^{18}\text{R}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{R}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{OR}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{NR}^{19}\text{R}^{20}$, $\text{C}(=\text{O})\text{R}^{18}$, $\text{C}(=\text{NOR}^{18})\text{R}^{19}$, $\text{C}(=\text{O})\text{OR}^{18}$, $\text{C}(=\text{O})\text{NR}^{18}\text{R}^{19}$, $\text{C}(=\text{O})\text{NR}^{18}\text{OR}^{19}$ or R^{21} ,
- 15

wherein,

- R^{18} , R^{19} , R^{20} and R^{21} independently is H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$ cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,
- 20

In still another embodiment,

- R^{17} and R^{24} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $\text{S}(=\text{O})\text{R}^{18}$, $\text{S}(=\text{O})_2\text{R}^{18}$, $\text{S}(=\text{O})_2\text{NR}^{18}\text{R}^{19}$, NO_2 , $\text{NR}^{18}\text{R}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{R}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{OR}^{19}$, $\text{NR}^{18}\text{C}(=\text{O})\text{NR}^{19}\text{R}^{20}$, $\text{C}(=\text{O})\text{R}^{18}$, $\text{C}(=\text{NOR}^{18})\text{R}^{19}$, $\text{C}(=\text{O})\text{OR}^{18}$, $\text{C}(=\text{O})\text{NR}^{18}\text{R}^{19}$, $\text{C}(=\text{O})\text{NR}^{18}\text{OR}^{19}$ or R^{21} ,
- 25

wherein,

- R^{18} , R^{19} , R^{20} and R^{21} independently is H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$ cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,
- 30

- 35 In still another embodiment,

R^{17} and R^{24} independently is H, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$,
 5 $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,
 wherein,
 R^{18} , R^{19} , R^{20} and R^{21} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8
 10 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,
 R^{17} and R^{24} independently is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the
 15 group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$,
 NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$,
 $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,
 wherein,
 R^{18} , R^{19} , R^{20} and R^{21} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8
 20 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,
 25 R^{17} and R^{24} independently is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} ,
 $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$,
 $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$,
 $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,
 30 wherein,
 R^{18} , R^{19} , R^{20} and R^{21} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8
 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,
 35

In still another embodiment,

R^{17} and R^{24} independently is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R^{17} and R^{24} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R^{17} and R^{24} independently is H, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^{18} and R^{19} may together form a 3-8 membered

heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

5 R^{17} and R^{24} independently is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

10 wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

15

In still another embodiment,

R^{17} and R^{24} independently is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

20

wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

25

In still another embodiment,

R^{17} and R^{24} independently is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

30

35

wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R^{17} and R^{24} independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,
wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, methyl, ethyl, propyl or butyl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R^{17} and R^{24} independently is H, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,
wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, methyl, ethyl, propyl or butyl and wherein R^{18} and R^{19} may together form a 3-8 membered heterocyclic ring or R^{18} and R^{20} may together form a 3-8 membered heterocyclic ring or R^{19} and R^{20} may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R^{17} and R^{24} independently is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$,

NO₂, NR¹⁸R¹⁹, NR¹⁸C(=O)R¹⁹, NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, C(=O)R¹⁸,
C(=NOR¹⁸)R¹⁹, C(=O)OR¹⁸, C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹ or R²¹,
wherein,

- 5 R¹⁸, R¹⁹, R²⁰ and R²¹ independently is H, methyl, ethyl, propyl or butyl and wherein
R¹⁸ and R¹⁹ may together form a 3-8 membered heterocyclic ring or R¹⁸ and R²⁰ may
together form a 3-8 membered heterocyclic ring or R¹⁹ and R²⁰ may together form a
3-8 membered heterocyclic ring,

In still another embodiment,

- 10 R¹⁷ and R²⁴ independently is H, phenyl or naphthyl optionally substituted with one or
more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹⁸,
S(=O)R¹⁸, S(=O)₂R¹⁸, S(=O)₂NR¹⁸R¹⁹, NO₂, NR¹⁸R¹⁹, NR¹⁸C(=O)R¹⁹,
NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, C(=O)R¹⁸, C(=NOR¹⁸)R¹⁹, C(=O)OR¹⁸,
C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹ or R²¹,
15 wherein,
R¹⁸, R¹⁹, R²⁰ and R²¹ independently is H, methyl, ethyl, propyl or butyl and wherein
R¹⁸ and R¹⁹ may together form a 3-8 membered heterocyclic ring or R¹⁸ and R²⁰ may
together form a 3-8 membered heterocyclic ring or R¹⁹ and R²⁰ may together form a
3-8 membered heterocyclic ring,

In still another embodiment,

- 20 R¹⁷ and R²⁴ independently is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl op-
tionally substituted with one or more substituents selected from the group consisting
of F, Cl, CN, CF₃, OR¹⁸, S(=O)R¹⁸, S(=O)₂R¹⁸, S(=O)₂NR¹⁸R¹⁹, NO₂, NR¹⁸R¹⁹,
25 NR¹⁸C(=O)R¹⁹, NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, C(=O)R¹⁸, C(=NOR¹⁸)R¹⁹,
C(=O)OR¹⁸, C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹ or R²¹,
wherein,
R¹⁸, R¹⁹, R²⁰ and R²¹ independently is H, methyl, ethyl, propyl or butyl and wherein
R¹⁸ and R¹⁹ may together form a 3-8 membered heterocyclic ring or R¹⁸ and R²⁰ may
30 together form a 3-8 membered heterocyclic ring or R¹⁹ and R²⁰ may together form a
3-8 membered heterocyclic ring,

In still another embodiment,

- 35 R¹⁷ and R²⁴ independently is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl,
cyclopentyl or cyclohexyl optionally substituted with one or more substituents se-

lected from the group consisting of F, Cl, CN, CF₃, OR¹⁸, S(=O)R¹⁸, S(=O)₂R¹⁸, S(=O)₂NR¹⁸R¹⁹, NO₂, NR¹⁸R¹⁹, NR¹⁸C(=O)R¹⁹, NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, C(=O)R¹⁸, C(=NOR¹⁸)R¹⁹, C(=O)OR¹⁸, C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹ or R²¹, wherein,

- 5 R¹⁸, R¹⁹, R²⁰ and R²¹ independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- 10 R¹⁷ and R²⁴ independently is aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹⁸, S(=O)R¹⁸, S(=O)₂R¹⁸, S(=O)₂NR¹⁸R¹⁹, NO₂, NR¹⁸R¹⁹, NR¹⁸C(=O)R¹⁹, NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, C(=O)R¹⁸, C(=NOR¹⁸)R¹⁹, C(=O)OR¹⁸, C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹ or R²¹, wherein,

- 15 R¹⁸, R¹⁹, R²⁰ and R²¹ independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- 20 R¹⁷ and R²⁴ independently is phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹⁸, S(=O)R¹⁸, S(=O)₂R¹⁸, S(=O)₂NR¹⁸R¹⁹, NO₂, NR¹⁸R¹⁹, NR¹⁸C(=O)R¹⁹, NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, C(=O)R¹⁸, C(=NOR¹⁸)R¹⁹, C(=O)OR¹⁸, C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹ or R²¹, wherein,

- 25 R¹⁸, R¹⁹, R²⁰ and R²¹ independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- 30 R¹⁷ and R²⁴ independently is phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF₃, OR¹⁸, S(=O)R¹⁸, S(=O)₂R¹⁸, S(=O)₂NR¹⁸R¹⁹, NO₂, NR¹⁸R¹⁹, NR¹⁸C(=O)R¹⁹, NR¹⁸C(=O)OR¹⁹, NR¹⁸C(=O)NR¹⁹R²⁰, C(=O)R¹⁸, C(=NOR¹⁸)R¹⁹, C(=O)OR¹⁸, C(=O)NR¹⁸R¹⁹, C(=O)NR¹⁸OR¹⁹ or R²¹, wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- 5 R^{17} and R^{24} independently is thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

10 wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- 15 R^{17} and R^{24} independently is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

20 wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

In still another embodiment,

- 25 R^{17} and R^{24} independently is aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,

30 wherein,

R^{18} , R^{19} , R^{20} and R^{21} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

In still another embodiment,

- R^{17} and R^{24} independently is phenyl, naphthyl, thienyl, furyl, pyridyl, quinoliny or isoquinoliny optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,
 5 wherein,
 R^{18} , R^{19} , R^{20} and R^{21} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinoliny or isoquinoliny,
- 10 In still another embodiment,
 R^{17} and R^{24} independently is phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$, $C(=O)OR^{18}$,
 15 $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,
 wherein,
 R^{18} , R^{19} , R^{20} and R^{21} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinoliny or isoquinoliny,
- 20 In still another embodiment,
 R^{17} and R^{24} independently is thienyl, furyl, pyridyl, quinoliny or isoquinoliny optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF_3 , OR^{18} , $S(=O)R^{18}$, $S(=O)_2R^{18}$, $S(=O)_2NR^{18}R^{19}$, NO_2 , $NR^{18}R^{19}$, $NR^{18}C(=O)R^{19}$, $NR^{18}C(=O)OR^{19}$, $NR^{18}C(=O)NR^{19}R^{20}$, $C(=O)R^{18}$, $C(=NOR^{18})R^{19}$,
 25 $C(=O)OR^{18}$, $C(=O)NR^{18}R^{19}$, $C(=O)NR^{18}OR^{19}$ or R^{21} ,
 wherein,
 R^{18} , R^{19} , R^{20} and R^{21} independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinoliny or isoquinoliny,
- 30 In still another embodiment,
 R^{17} and R^{24} independently is H, C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl or heteroaryl
- 35 In still another embodiment,
 R^{17} and R^{24} independently is H,

In still another embodiment,

R^{17} and R^{24} independently is C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl or C_3 - C_7 cycloheteroalkyl,

5 In still another embodiment,

R^{17} and R^{24} independently is methyl, ethyl, propyl or butyl

in still another preferred embodiment

R^{17} and R^{24} independently is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl

10

in still another preferred embodiment

R^{17} and R^{24} independently is aziridiny, pyrrolidiny, piperidiny or morpholiny

In still another embodiment,

15 R^{17} and R^{24} independently is aryl or heteroaryl

In still another embodiment,


R^{17} and R^{24} independently is phenyl or naphthyl

20 In still another embodiment,

R^{17} and R^{24} independently is thienyl, furyl, pyridyl, quinoliny or isoquinolyl

Experiments

25 General Procedure 1: Synthesis of benzoic acid derivatives for building blocks:




The benzoic acid derivative (1 mmol) was dissolved in THF (5 mL) and pyridine (3 mmol). The mixture was cooled to 0°C and treated with an acid chloride (1.2 mmol).

30 The cooling bath was removed and the reaction mixture was stirred for 1 hour at rt. Toluene (10 mL) was added and the solution was evaporated *in vacuo*. The crude

was redissolved in EtOAc (10 mL), washed with water and brine. The organic phase was dried over MgSO_4 and evaporated *in vacuo*. The pure product was obtained by silica gel purification using a gradient of heptane to EtOAc as eluent.

Example 1 (General procedure 1, wherein $\text{Z}=\text{S}$, $\text{R}'=\text{H}$, and $\text{R}=\text{CH}_3$)


5 4-Acetylsulfanyl-benzoic acid



Yield = 70%; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): 8.00 (d, 2H); 7.55 (d, 2H); 2.46 (s, 3H).

Example 2 (General procedure 1, wherein $\text{Z}=\text{S}$, $\text{R}'=\text{H}$, and $\text{R}=\text{CH}_2\text{CH}_3$)


10 4-Propionylsulfanyl-benzoic acid



Yield = 85%; $^1\text{H-NMR}$ (CDCl_3): 8.12 (d, 2H); 7.58 (d, 2H); 2.76 (q, 2H); 1.28 (t, 3H).

Example 3 (General procedure 1, wherein $\text{Z}=\text{S}$, $\text{R}'=\text{H}$, and $\text{R}=(\text{CH}_2)_2\text{CH}_3$)


15 4-Butyrylsulfanyl-benzoic acid



Yield = 98%: ^1H NMR (CDCl_3): 8.15 (d, 2H); 7.56 (d, 2H); 2.70 (t, 2H); 1.81 (sextet, 2H); 1.04 (t, 3H) .

5 **Example 4** (General procedure 1, wherein $\text{Z}=\text{S}$, $\text{R}'=\text{H}$, and $\text{R}=(\text{CH}_2)_2\text{CHCH}_2$)

4-Pent-4-enoylsulfanylbenzoic acid




Yield = 71%: ^1H -NMR (CDCl_3): 8.15 (d, 2H); 7.55 (d, 2H); 5.85 (m, 1H); 5.11 (dd, 2H); 2.82 (t, 2H); 2.47 (q, 2H).

10


Example 5 (General procedure, wherein $\text{Z}=\text{O}$, $\text{R}'=\text{Cl}$, and $\text{R}=\text{CH}_3$)

4-Acetoxy-3-chloro-benzoic acid



Yield = 95%: ^1H nmr (CDCl_3): 8.20 (d, 1H); 8.05 (dd, 1H), 7.25 (d, 1H); 2.40 (s, 3H).


15

General Procedure 2: Synthesis of nicotinic acid derivative for building blocks:


5 The nicotinic acid derivative (6.44 mmol) was dissolved in THF (10 mL) and triethyl-
amine (5 mL). The mixture was cooled to 0°C and treated with an acid chloride
(12.88 mmol). The cooling bath was removed and the reaction mixture was stirred
overnight at rt. After removal of the solvents, toluene (10 mL) was added to the
crude and evaporated *in vacuo*. The pure product was obtained by silica gel purifica-
10 tion using a gradient starting from dichloromethane going to 2% methanol in di-
chloromethane as eluent.

Example 6 (General procedure 2, wherein Z=S, R'=H, and R= CH₃)

2-Acetylsulfanyl-nicotinic acid



15 Yield = 5%: ¹H-NMR (CDCl₃): 8.76 (dd, 1H); 8.64 (dd, 1H); 7.40 (dd, 1H); 2.79 (s,
3H).

General Procedure 3: Preparation of building blocks by loading a Carrier-Functional entity ensemble onto an oligonucleotide comprising an amino group:

25 μ L of a 150 mM benzoic acid derivative in DMF was mixed with 25 μ L of a 150 mM solution of EDC in DMF. The mixture was left for 30 min at 25°C. 50 μ L of an aminooligo (10 nmol) in 100 mM HEPES buffer pH 7.5 was added and the reaction mixture was left for 20 min at 25°C. The excess building block was removed by extraction with EtOAc (500 μ L) and remaining EtOAc was removed *in vacuo* by spinning 10 min in a speedvac. The aminooligo loaded with the benzoic acid derivative was ethanol precipitated twice using NH_4OAc and analysed by electron spray mass spectrometry (ES-MS).

10 Aminooligo's used:

A: 5'-XTTTTTTTTTTTTTTTTACGACTACGTTTCAGGCAAGTB

B: 5'-XTTTTTTTTTTTTTTTTTTTTACGACTACGTTTCAGGCAAGTB

C: 5'-XTTTTTTTTTTTTTTTTTTTTTTACGACTACGTTTCAGGCAAGTB

15 D: 5'-BGACCTGTCGAGCATCCAGCZ

E: 5'-BGCATCCATCGY

X = 5' amino C6 (Glen# 10-1906-90)


Y = C2 amino dT phosphate (Glen# 10-1037-90)

20 Z = C6 amino dT phosphate (Glen# 10-1039)

B = Biotin (Glen # 10-1953-95)

Example 7 (General procedure (3))

Oligo A loaded with compound of Example 1




25

MS (calc., M-1) = 11.560,87 ;MS (found) = 11.557,89

Example 8 (General procedure (3))


Oligo B loaded with compound of Example 1



MS (calc., M-1) = 13.081,87; MS (found) = 13.079,01

Example 9 (General procedure (3))


Oligo C loaded with compound of Example 1



MS (calc., M-1) = 14.602,86; MS (found) = 14.599,66

Example 10 (General procedure (3))


Oligo D loaded with compound of Example 1



MS (calc., M-1) = 6892,85; MS (found) = 6893,29

Example 11 (General procedure (3))


Oligo E loaded with compound of Example 1



MS (calc., M-1) = 4052,05; MS (found) = 4067,49¹


Example 12 (General procedure (3))

Oligo E loaded with compound of Example 5



MS (calc., M-1) = 4069,84; MS (found) = 4070,20

10 General Procedure 4: Preparation of building blocks by step wise loading of a Carrier and a Functional Entity onto an oligonucleotide containing a nucleotide derivative comprising an amino group:



40 μ L of a 20 mM SPDP solution in DMSO was mixed with an aminooligo (5 nmol). 200 mM HEPES buffer pH 7.5 was added (80 μ L) and water to a final volume of 160 μ L. the reaction mixture was left for 2 hours at 30°C. The excess building block was removed by extraction with EtOAc (500 μ L). Remaining EtOAc was removed *in*

¹ The difference observed in the calculated and found MS of around 16 is probably due to an oxidation of the sulphur atom of the biotin moiety

vacuo by spinning 10 min in a speedvac. The SPDP activated aminooligo was purified using a micro bio-spin column (equilibrated with 200 mM HEPES buffer pH 7.5). 10 μ L of a 50 mM thio acid derivate solution in DMSO was added to the purified SPDP activated aminooligo solution and the reaction mixture was left for 30 min at 20°C. The building block loaded aminooligo was ethanol precipitated twice using NH_4OAc and analysed by electron spray mass spectrometry (ES-MS).


Aminooligo used:

A2: 5'- GACCTGTCGAGCATCCAGCTTCATGGGAATTCCTCGTCCACAATGZ

Z = Amino-Modifier C6 dT phosphate (Glen# 10-1039-)

Example 13 (General procedure (4))

Oligo A2 loaded with thiobenzoic acid




MS (calc., M-1) = 14518,76; MS (found) = 14516,78

Example 14: Loading of a trisamine scaffold on an oligonucleotide containing a nucleotide derivative comprising an amino group:

A hexameric scaffold peptide with the sequence, CysPhePheLysLysLys, was synthesised by standard solid-phase Fmoc peptide chemistry. The scaffold peptide comprises a -SH group on the cystein side chain, said -SH group being used for coupling the scaffold peptide to a amine-bearing oligonucleotide serving as anti-codon and linker. Each of the three lysin moieties comprises an amino group in the side chain. The amine groups are used as reactive groups for the formation of a connection to functional entities emanating from building blocks.

The N-terminus of the peptide was acetylated and the C-terminus was initially capped as an amide to avoid any participation in the reactions to follow and subsequently purified by reverse phase-HPLC. The scaffold peptide was covalently attached to DNA oligonucleotide using the scheme shown schematically below. For illustrative purposes, the scaffold is indicated as HS—Scaffold




5 nmol of oligodeoxynucleotide F: 5'-XTCGTAACGACTGAATGACGT where X = 5' amino C6 (Glen# 10-1906-90) in 100 mM Hepes-OH pH 7.5 is incubated with 20 mM Succinimidyl-propyl-2-dithiopyridyl (SPDP, Molecular probes) dissolved in DMSO for 3 hours at 25 °C. Excess SPDP is removed by triple extraction using 5 volumes of ethylacetate. The sample is further purified using a Bio-rad Microspin 6 column equilibrated in H₂O.

The oligonucleotide-scaffold conjugate is synthesised by incubating 1 µmol hexapeptide with 5 nmol SPDP activated oligonucleotide in 100 mM Hepes-OH pH 7.5 for 2 hours at 25 °C. Excess peptide is removed by double sodium-acetate/ethanol precipitation of the scaffold-DNA complex according to standard procedure. The loading was verified by Electrospray Mass Spectrometry (ES-MS).

Loading of trisamine scaffold on oligo F: MS (calc., M-1) = 7247.45 MS (found) = 7244.80

Example 15: Transfer of a functional entity from a building block to a scaffold:



A template oligo G: 5'-ACGTCATTACGTCGTTACGAACGATGGATGCTCCAGG TCGC (1 nmol) was mixed with scaffold oligo F (1.5 nmol) in MES-buffer (20 µL of a 100 mM MES, pH=6) and water (added to a final volume of 100 µL). Scaffold oligo F was annealed to the template by heating to 80 °C and cooled (-2 °C/ 10 second) to room temperature and functional entity oligo E (Example 11) (1.5 nmol) was added. The mixture was left o/n at room temperature. The oligo complex was attached to

streptavidine by addition of streptavidine sepharose beads (50 μ L, prewashed with 2x1 mL 100 mM MES buffer, pH=6). The beads were washed with water (4 x 200 μ L). Oligo F was separated from the streptavidine bound complex by addition of water (200 μ L) followed by heating to 80 °C for 5 minute. The beads were filtered off and the water was evaporated. Oligo F was redissolved in water and building block transfer verified by electron spray mass spectrometry (ES-MS).

Transfer of acetyl to trisamine scaffold oligo F from example I attached to oligo E:
MS (calc.) = 7289.49; MS (found) = 7286.58

Section 3: Transfer efficiencies of functional entities from building blocks to amine scaffolds

Carrier coupled functional entities were loaded onto oligos (oligonucleotides) containing a nucleotide derivative comprising an amino group (General procedure 5) or a nucleotide derivative comprising a thiol (General procedure 6) and the transfer was conducted to a scaffold oligo with a nucleotide derivative comprising an amino group. Transfer efficiencies were analyzed by ES-MS (electrospray mass spectroscopy) (General procedure 7).

General Procedure 5: Loading of a carrier coupled functional entity onto an amino oligo:

25 μ L 100 mM carrier coupled functional entity dissolved in DMF (dimethyl formamide) was mixed with 25 μ L 100 mM EDC (1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride) in DMF for 30 minutes at 25° C. The mixture was added to 50 μ L amino oligo in H₂O with 100 mM HEPES (2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-ethanesulfonic acid) pH 7.5 and the reaction was allowed to proceed for 20 minutes at 25° C. Unreacted carrier coupled functional entity was removed by extraction with 500 μ L EtOAc (ethyl acetate), and the oligo was purified by gel filtration through a microspin column equilibrated with 100 mM MES (2-(N-morpholino) ethanesulfonic acid) pH 6.0.


Oligonucleotide used:

Oligo G: 5'-GCGACCTGGAGCATCCATCGY

Y = Amino-Modifier C2 dT phosphate (Glen# 10-1037)

Example 16 (General procedure 5, using compound of Example 5 as carrier coupled functional entity)


- 5 Carrier coupled functional entity: 4-Acetoxy-3-chloro-benzoic acid



- 10 Mass: 6738.23 (observed using ES-MS), 6738.31 (calculated) (The carrier coupled functional entity oligo is hydrolyzed in the mass spectrometer during analysis).

Example 17 (General procedure 5, using compound of example 1 as carrier coupled functional entity)


Carrier coupled functional entity: 4-Acetylsulfanyl-benzoic acid



- 15 Mass: 6718.48 (observed using ES-MS), 6719.48 (calculated) (The carrier coupled functional entity oligo is hydrolyzed in the mass spectrometer during analysis).

Example 18 (General procedure 1, wherein Z=O, R'=NO₂, and R=CH₃ and general procedure 5)

- 20 Carrier coupled functional entity: 4-Acetoxy-3-nitro-benzoic acid



Mass: 6748.31 (observed using ES-MS), 6748.42 (calculated) (The carrier coupled functional entity oligo is hydrolyzed in the mass spectrometer during analysis).


General Procedure 6: Loading of a carrier coupled functional entity onto a thiol oligo:

- 5 10 nmol thiol oligo was lyophilized and redissolved in 50 μ l H₂O with 100 mM dithiothreitol and 100 mM sodium phosphate pH 8.0 and incubated at 37 °C for 1 hour. The reduced oligo was purified using a microspin column equilibrated with HEPES (100 mM, pH 7.5). Then 100 mM NHM (N-hydroxymaleimide) in HEPES (100 mM, pH 7.5) was added to the thiol oligo and the mixture was incubated at 25°C for 2
- 10 hours. The resulting NHS (N-hydroxysuccinimide)-oligo was purified using a microspin column equilibrated with H₂O. 1 nmol NHS-oligo was lyophilized and redissolved in 10 μ l 100 mM MES, pH 6. 50 μ l carrier coupled functional entity (100 mM) in dimethyl formamide was activated with 50 μ l 100 mM EDC in DMF for 30 min at 25 °C. 10 μ l of the EDC-activated carrier coupled functional entity was mixed with
- 15 the NHS-oligo and incubated for 5 min at 25 °C. 30 μ l 100 mM MES pH 6 was added and following an extraction with 500 μ l EtOAc the oligo was purified using a microspin column equilibrated with 100 mM MES pH6.

Oligo H: 5'-GCGACCTGGAGCATCCATCGTX


- 20 X = Thiol-Modifier C6 S-S (Glen# 10-1936)

Example 19 (General procedure 6)



- 25 Mass "X": 6723.21 (observed using ES-MS), 6723.52 (calculated) (Compound "Z" is hydrolyzed to compound "X" in the mass spectrometer during analysis).

General procedure 7: Transfer of functional entity from a carrier oligo to a scaffold oligo.




Scaffold oligo I: 5'- ZACGATGGATGCTCCAGGTCGC

Z = 5' Amino-modifier C6 (Glen Research cat. # 10-1906)

- 10 A carrier coupled functional entity oligo (Examples 16, 17, 18, 19) (250 pmol) was added to a scaffold oligo I (200 pmol) in 50 μ l 100 mM MES, pH 6. The mixture was incubated overnight at 25 $^{\circ}$ C. Subsequently, the mixture was purified by gel filtration using a microspin column equilibrated with H₂O and transfer of the functional entity was verified by electron spray mass spectrometry (ES-MS). Transfer efficiencies are expressed in percent and were calculated by dividing the abundance of scaffold oligo carrying transferred functional entities to total abundance of scaffold oligos (with and without transferred functional entities).
- 15

Example 20 (General procedure 7):

20



Mass ("X"): 6624.70 (observed), 6625.42 (calculated). Abundance: 73.16 (arbitrary units)





25 Mass ("Y"): 6666.09 (observed), 6667.46 (calculated). Abundance: 26.15 (arbitrary units)

Mass ("Z"): 6738.01 (observed), 6738.31 (calculated) (carrier coupled functional entity oligos are hydrolyzed in the mass spectrometer during analysis).

Transfer efficiency calculated as: $26.15 / (26.15 + 73.16) = 0.2633 \sim 26 \%$

Transfer efficiencies:

5

Scaffold oligo	Building block oligo			
	Example 16	Example 17	Example 18	Example 19
				
	26	32	>60	58

Example 21: Stability of building block oligonucleotides during storage and handling

- 10 Carrier coupled functional entities were loaded onto oligonucleotides containing a nucleotide derivative comprising an amino group (General Procedure 7). The resulting carrier coupled functional entity oligos were either mixed immediately with scaffold oligo I at 25°C (condition 1) or subjected to different conditions before mixing: (condition 2) -80°C for 14 days, (condition 3) 25°C for 1 hour. For condition 4
- 15 the scaffold oligo and the building block oligo were heated to 80°C for 30 seconds, mixed, and then cooled to 25°C (-5°C / minute). The functional entity of the building block oligo was transferred to a scaffold oligo by incubation at 25°C overnight and analyzed by ES-MS (General procedure 3).
- 20 Transfer efficiencies (in percent) in reactions involving the same building block were normalized to facilitate comparison, e.g. the observed transfer efficiency when scaffold oligo was mixed with building block oligo immediately after production was set to 100:

Condition	Description	Ex. 16	Ex. 17	Ex. 18	Ex. 19
1	Immediate mixing	100	100	100	100
2	-80°C for 14 days	96	97	89	92
3	25°C for 1 hour	97	98	93	71
4	From 80°C to 25°C	106	105	87	60 ⁵

10 The results indicate that all the building blocks may be stored in a freezer at -80°C for several weeks without losing significant reactivity. Under practical handling conditions at room temperature the NHS ester of example 19, which is not according to the invention, loses a considerable amount of reactivity. The tendency of spontaneous hydrolysis of the building block according to example 18 is reinforced under the condition simulating an actual experiment (condition 4), while the building blocks of example 16 to 18 have an acceptable stability or even a slightly increased activity. Activities above 100 observed under condition 4 might be due to experimental variation or facilitation of annealing of the carrier coupled functional entity oligo and scaffold oligo at elevated temperatures.

20 **Example 22: Preparation of Building blocks.**

The following oligo containing a nucleobase modified with an amino group was synthesised, using the conventional phosphoramidite approach:



25 N: 5'-ZGT AAC ACC TGT GTA AGC TGC CTG TCA GTC GGT ACT GAC CTG
TCG AGC ATC CAG CT

30 Z depicts the nucleobase modified with an aminogroup, incorporated using the commercially available amino modifier C6 dT phosphoramidite (10-1039-90 from Glen research)

The loading with a functional entity proceeds using the general method:
An amino oligo (3 pmol) was mixed with a phosphate buffer (3 uL of a 0.1 M solution, pH=6) and NaBH₃CN (3 uL of a 1 M solution in MeOH). A chemical entity com-

prising the functional entity (3 μ L of a 1 M solution in MeOH) was added and the mixture was left o/n at room temperature. The product formation was analysed by PAGE gel.

- 5 Exemplary chemical entities are 4-acetoxybenzaldehyde (24,260-8 from Sigma-Aldrich),



Propionic acid 4-formyl-phenyl ester

, and Butanoic acid 4-formyl-phenyl ester

- 10 Figure 5 shows a PAGE analysis of the loading of an oligo with butanoic acid 4-formyl-phenyl ester. Lane 1 shows the reference amino oligo (N). Lane 2 show the amino oligo (N) after loading with a the chemical entity comprising the functional entity, and Lane 3 shows removal of the functional entity, attached in lane 2, by treatment with pH=11 for 1 hour.

- 15 The above examples are intended to help illustrate the invention, and are not intended to, nor should they be construed to, limit the scope of the invention. Indeed, various modifications of the invention and many further embodiments thereof, in addition to those shown and described herein, will become apparent to those skilled in the art from the full content of this document, including the examples shown
- 20 above and the references to the scientific a patent literature cited herein. It should further be appreciated that the contents of those cited references are incorporated herein by reference to help illustrate the state of the art. The examples above contain important additional information that can be adapted to the practice of this invention in its various embodiments and the equivalents thereof.

25

Abbreviations

DCC	N,N'-Dicyclohexylcarbodiimide
DhbtOH	3,4-dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine
DIC	Diisopropylcarbodiimide
DIEA	Diethylisopropylamin
DMAP	4-Dimethylaminopyridine
DNA	Deoxyribosenucleic Acid
EDC	1-Ethyl-3-(3'-dimethylaminopropyl)carbodiimide·HCl
HATU	2-(1 <i>H</i> -7-Azabenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HBTU	2-(1 <i>H</i> -Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HOAt	N-Hydroxy-7-azabenzotriazole
HOBt	N-Hydroxybenzotriazole
LNA	Locked Nucleic Acid
NHS	N-hydroxysuccinimid
OTf	Trifluoromethylsulfonate
OTs	Toluenesulfonate
PNA	Peptide Nucleic Acid
PyBoP	Benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate
PyBroP	Bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
RNA	Ribonucleic acid
TBTU	2-(1 <i>H</i> -Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate
TEA	Triethylamine
RP-HPLC	Reverse Phase High Performance Liquid Chromatography
TBDMS-Cl	<i>Tert</i> -Butyldimethylsilylchloride
5-Iodo-dU	5-iodo-deoxyriboseuracil
TLC	Thin layer chromatography
(Boc) ₂ O	Boc anhydride, di- <i>tert</i> -butyl dicarbonate
TBAF	Tetrabutylammonium fluoride
SPDP	Succinimidyl-propyl-2-dithiopyridyl

Claims

1. A building block of the general formula


Complementing Element – Linker – Carrier – C-F-connecting group – Functional entity precursor

capable of transferring a functional entity to a recipient reactive group, wherein

Complementing Element is a group identifying the functional entity precursor,

Linker is a chemical moiety comprising a **Spacer** and a **S-C-connecting group**, wherein the **Spacer** is a valence bond or a group distancing the functional entity precursor to be transferred from the complementing element and the S-C-connecting group connects the spacer with the Carrier,

Carrier is selected among the groups



wherein the Linker attaches to the Carrier through Y and

W = CH or N

$R^2 = -H, -\text{Halogen}, -\text{NO}_2, -\text{CN}, -\text{C}(\text{Halogen})_3, -\text{C}(\text{O})\text{R}^3, -\text{C}(\text{O})\text{NHR}^3, \text{C}(\text{O})\text{NR}^3_2,$
 $-\text{NC}(\text{O})\text{R}^3, -\text{S}(\text{O})_2\text{NHR}^3, -\text{S}(\text{O})_2\text{NR}^3_2, -\text{S}(\text{O})_2\text{R}^3, -\text{P}(\text{O})_2\text{R}^3, -\text{P}(\text{O})-\text{R}^3, -\text{S}(\text{O})-\text{R}^3,$
 $\text{P}(\text{O})-\text{OR}^3, -\text{S}(\text{O})-\text{OR}^3, -\text{N}^+\text{R}^3_3,$ wherein p is an integer of 0 to 3, $\text{R}^3 = \text{H}, \text{C}_1\text{-C}_6 \text{ alkyl},$
 $\text{C}_1\text{-C}_6 \text{ alkenyl}, \text{C}_1\text{-C}_6 \text{ alkynyl}, \text{or aryl},$ and Halogen is F, Cl, Br, or I,
 Y = absent, $\text{C}_1\text{-C}_6 \text{ Alkylene}, \text{C}_1\text{-C}_6 \text{ Alkenylene}, \text{C}_1\text{-C}_6 \text{ Alkynylene}, \text{Arylene}, \text{Heteroarylene}, \text{Carbonyl}, \text{or } -\text{SO}_2\text{CH}_2-$

25 **C-F-connecting group** is $-\text{Z}-\overset{\text{V}}{\underset{\text{X}}{\text{C}}}-$ or $-\overset{\text{V}}{\underset{\text{X}}{\text{C}}}-$ where the carrier is connected to the left hand side of the formulae and
 $\text{X} = -\text{C}-, -\text{S}-, -\text{P}-, -\text{S}(\text{O})-, \text{or } -\text{P}(\text{O})-,$
 $\text{V} = \text{O}, \text{S}, \text{NH}, \text{or } \text{N-C}_1\text{-C}_6 \text{ alkyl}, \text{and}$
 $\text{Z} = \text{O}, \text{S}; \text{and}$

30 **Functional entity precursor** is H or selected among the group consisting of a $\text{C}_1\text{-C}_6 \text{ alkyl}, \text{C}_2\text{-C}_6 \text{ alkenyl}, \text{C}_2\text{-C}_6 \text{ alkynyl}, \text{C}_4\text{-C}_8 \text{ alkadienyl}, \text{C}_3\text{-C}_7 \text{ cycloalkyl}, \text{C}_3\text{-C}_7 \text{ cycloheteroalkyl}, \text{aryl}, \text{and heteroaryl},$ said group being substituted with 0-3 R^4 , 0-3

R^5 and 0-3 R^9 , or selected among the group consisting of C_1 - C_3 alkylene- NR^4 , C_1 - C_3 alkylene- $NR^4C(O)R^8$, C_1 - C_3 alkylene- $NR^4C(O)OR^8$, C_1 - C_2 alkylene- $O-NR^4$, C_1 - C_2 alkylene- $O-NR^4C(O)R^8$, and C_1 - C_2 alkylene- $O-NR^4C(O)OR^8$ substituted with 0-3 R^9 .

where R^4 is H or selected independently among the group consisting of C_1 - C_8

- 5 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, C_3 - C_7 cycloalkyl, C_3 - C_7 cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R^9 and


R^5 is selected independently from $-N_3$, $-CNO$, $-C(NOH)NH_2$, $-NHOH$, $-NHNHR^6$, $-C(O)R^6$, $-SnR^6_3$, $-B(OR^6)_2$, $-P(O)(OR^6)_2$ or the group consisting of C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, C_4 - C_8 alkadienyl said group being substituted with 0-2 R^7 ,

- 10 where R^6 is selected independently from H, C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, aryl or C_1 - C_8 alkylene-aryl substituted with 0-5 halogen atoms selected from $-F$, $-Cl$, $-Br$, and $-I$; and R^7 is independently selected from $-NO_2$, $-COOR^6$, $-COR^6$, $-CN$, $-OSiR^6_3$, $-OR^6$ and $-NR^6_2$.

- R^8 is H, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, C_3 - C_7 cycloalkyl, aryl or C_1 - C_8 alkylene-aryl substituted with 0-3 substituents independently selected from $-F$, $-Cl$, $-NO_2$, $-R^3$, $-OR^3$, $-SiR^3_3$

R^9 is $=O$, $-F$, $-Cl$, $-Br$, $-I$, $-CN$, $-NO_2$, $-OR^6$, $-NR^6_2$, $-NR^6-C(O)R^8$, $-NR^6-C(O)OR^8$, $-SR^6$, $-S(O)R^6$, $-S(O)_2R^6$, $-COOR^6$, $-C(O)NR^6_2$ and $-S(O)_2NR^6_2$.

- 20 2. The compound according to claim 1, wherein the **Spacer** is a valence bond, C_1 - C_8 alkylene-A-, C_1 - C_8 alkenylene-A-, C_2 - C_8 alkynylene-A-, or



said spacer optionally being connected through A to a linker selected from




- 25 $-(CH_2)_n-S-S-(CH_2)_m-B-$

where A is $-C(O)NR^1$, $-NR^1$, $-O$, $-S$, or $-C(O)-O$; B is $-O$, $-S$, $-NR^1$ or $-$


$C(O)NR^1$ and connects to S-C-connecting group; R^1 is selected independently from H, C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, C_1 - C_8 alkylene-aryl, or aryl substituted with 0-5 halogen atoms selected from $-F$, $-Cl$, $-Br$ and $-I$; and n and m independently are integers

- 30 ranging from 1 to 10.

3. The compound according to claim 1, wherein the **Spacer** is C₁-C₈ alkylene-A-, C₁-C₈ alkenylene-A-, C₂-C₈ alkynylene-A-, or




said spacer optionally being connected through A to a moiety selected from




where A is -C(O)NR¹-, or -S-; B is -S-, -NR¹- or -C(O)NR¹- and connects to S-C-connecting group; R¹ is selected independently from H, C₁-C₆ alkyl, C₁-C₆ alkenylene-aryl, or aryl; and n and m independently are integers ranging from 1 to 6.

10

4. The compound according to claim 1, wherein **Spacer** is -A-, a group C₁-C₈ alkenylene-A-, C₂-C₈ alkenylene-A-, or C₂-C₈ alkynylene-A- optionally substituted with 1 to 3 hydroxy groups, or



15 said spacer being connected through A to a linker selected from




where A is a valence bond, -NR¹⁰-, -C(O)NR¹⁰-, -NR¹⁰-C(O)-, -O-, -S-, -C(O)-O- or -OP(=O)(O⁻)-O-; B is a valence bond, -O-, -S-, -NR¹⁰-, -C(O)- or -C(O)NR¹⁰- and connects to S-C-connecting group; R¹⁰ is selected independently from H, C₁-C₆ al-



20

kyl, C₃-C₇ cycloalkyl, aryl, C₁-C₆ alkylene-aryl, $-(CH_2)_4-O-(CH_2)_4-O-$ with a subscript n, or $-(CH_2)_4-N(G)-(CH_2)_4-N(G)-$ with a subscript n; G is H or C₁-C₆ alkyl; and n and m independently are integers ranging from 1 to 10.


25 5. A compound according to claim 4, wherein the **spacer** is C₂-C₈ alkenylene-A-, said spacer being connected through A to a moiety selected from




where A is a valence bond, $-\text{C}(\text{O})\text{NR}^{10}$ -, $-\text{NR}^{10}-\text{C}(\text{O})$ -, $-\text{S}$ -, $-\text{C}(\text{O})-\text{O}$ - or $-\text{OP}(=\text{O})(\text{O}^-)$ -; B is a valence bond, $-\text{S}$ -, $-\text{NR}^{10}$ -, or $-\text{C}(\text{O})$ - and connects to S-C-connecting group; n and m independently are integers ranging from 1 to 10 and

R¹⁰ is selected independently from H,  or , wherein G is H or C₁-C₆ alkyl; and the spacer is connected to the complementing element through a nucleobase.



6. A compound according to claim 4, wherein the spacer is -A-,



10 said spacer being connected through A to a moiety selected from





where A is a valence bond, -NR¹⁰-C(O)-, -O-, or -S-; B is a valence bond, -S-, -NR¹⁰-, or -C(O)- and connects to S-C-connecting group;
n and m independently are integers ranging from 1 to 10 and

15 R¹⁰ is selected independently from H,  or , wherein G is H or C₁-C₈ alkyl; and the spacer is connected to the complementing element via a phosphorus group.

7. A compound according to claim 6, wherein the phosphorus group is a phosphate or thiophosphate group attached to a 3' or 5' end of a complementing element.

8. A compound according to claims 1 to 7, wherein the **S-C-connecting group** is a valence bond, -NH-C(=O)- , $\text{-NH-C(=O)-C}_1\text{-C}_6$ alkylene-, -S-S- , $\text{-S-S-C}_1\text{-C}_6$ alkylene-, $\text{-C}_1\text{-C}_6$ alkylene-S-S -, $\text{-C(=O)-NH-(C}_1\text{-C}_6\text{ alkylene)-}$,






—NH-C(=O)-Arylene-C(R¹⁰)₂-NH-C(=O)-, -C(=O)-, -C(=O)-C₁-C₆ alkylene- or -C(=O)-Arylene-C(R¹⁰)₂-NR¹⁰-C(=O)-, where the right hand side of the formulae connects to the carrier.

5

9. A compound according to claims 1 to 8, wherein the **S-C-connecting group** is a valence bond, -NH-C(=O)-, -NH-C(=O)-C₁-C₆ alkylene-, -S-S-, -S-S-C₁-C₆ alkylene-, -C(=O)-NH-(C₁-C₆ alkylene)-,



10

-NH-C(=O)-Arylene-C(R¹⁰)₂-NH-C(=O)-, where the right hand side of the formulae connects to the carrier.

10. A compound according to claims 1 to 9, wherein the **S-C-connecting group** is -S-S-, -C₁-C₆ alkylene-S-S-, -C(=O)-NH-(C₁-C₆ alkylene)-, -C(=O)-, or -C(=O)-Arylene-C(R¹⁰)₂-NR¹⁰-C(=O)-, where the right hand side of the formulae connects to the carrier.

15


11. A compound according to claims 1 to 10, wherein the **S-C-connecting group** is -S-S-, -C(=O)-, or -C(=O)-Arylene-C(R¹⁰)₂-NR¹⁰-C(=O)-, where the right hand side of the formulae connects to the carrier.

20

12. The compound according to any of the claims 1 to 11, wherein the **S-C-connecting group** is a valence bond, -NH-C(=O)-, -S-S-, or -C(=O)-NH-, where the right hand side of the formulae connects to the carrier.


25

13. A compound according to claims 1 to 12, wherein the carrier is



and attaches to the linker through Y, and W, Y, R², and p are as defined in claim 1.

14. A compound according to claims 1 to 13, wherein the carrier is



5

and attaches to the linker through Y and

W = CH

10 R² = -H, halogen, -NO₂, -CN, -C(Halogen)₃, -C(O)R³, -C(O)NHR³, C(O)NR³,
-S(O)₂NHR³, -S(O)₂NR³, -S(O)₂R³, -N⁺R³, wherein halogen is selected from the
group consisting of -Cl, -F, -Br, and -I, p is an integer of 0 to 3, and R³ = H, C₁-C₈
alkyl, or aryl,

Y = absent, C₁-C₈ Alkylene, or carbonyl.

15 15. A compound according to any of the claims 1 to 14, wherein the C-F-connecting

group is $\text{---Z---}\overset{\text{V}}{\underset{\text{X}}{\text{C}}}$, in which

Z = O, S

X = -C-, and

V = O.

20

16. A compound according to any of the claims 1 to 15, wherein Complementing
element is a nucleic acid.

25 17. A compound according to any of the claims 1 to 16, wherein Complementing
element is a sequence of nucleotides selected from the group of DNA, RNA, LNA
PNA, morpholino derivatives, or combinations thereof.

18. A compound according to any of the claims 1 to 17, wherein the **Functional
entity precursor** is H or selected among the group consisting of a C₁-C₈ alkyl,
30 C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroal-

kyl, aryl, and heteroaryl, said group being substituted with 0-3 R⁵ and 0-3 R⁹, or selected among the group consisting of C₁-C₃ alkylene-NR⁴₂, C₁-C₃ alkylene-NR⁴C(O)R⁸, C₁-C₃ alkylene-NR⁴C(O)OR⁸, C₁-C₂ alkylene-O-NR⁴₂, C₁-C₂ alkylene-O-NR⁴C(O)R⁸, and C₁-C₂ alkylene-O-NR⁴C(O)OR⁸ substituted with 0-3 R⁹.

5

19. A compound according to claims 1 to 18, wherein the **Functional entity precursor** is H or selected among the group consisting of C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₄-C₈ alkadienyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloheteroalkyl, aryl, and heteroaryl, said group being substituted with 0-3 R⁵ and 0-3 R⁹.

10

20. A compound according to any of the claims 1 to 19, wherein **Functional entity precursor** is selected among the group consisting of C₁-C₃ alkylene-NR⁴₂, C₁-C₃ alkylene-NR⁴C(O)R⁸, C₁-C₃ alkylene-NR⁴C(O)OR⁸, C₁-C₂ alkylene-O-NR⁴₂, C₁-C₂ alkylene-O-NR⁴C(O)R⁸, and C₁-C₂ alkylene-O-NR⁴C(O)OR⁸ substituted with 0-3 R⁹.

15

21. A library of compounds according to any of the claims 1 to 20, wherein each different member of the library comprises a complementing element having a unique sequence of nucleotides, which identifies the functional entity.

20

22. A method for transferring a functional entity to a recipient reactive group, comprising the steps of

providing one or more building blocks according to any of the claims 1 to 20,

contacting the one or more building blocks with a corresponding encoding element associated with a recipient reactive group under conditions which allow for a recognition between the one or more complementing elements and the encoding elements, said contacting being performed prior to, simultaneously with, or subsequent to a transfer of the functional entity to the recipient reactive group.

25

30

23. The method according to claim 22, wherein the encoding element comprises one or more encoding sequences comprised of 1 to 50 nucleotides and the one or more complementing elements comprise a sequence of nucleotides complementary to one or more of the encoding sequences.

24. The method of claims 22 or 23, wherein the recipient reactive group is an amine group, which may be part of a chemical scaffold, and the linkage between the functional entity precursor and the scaffold is of the general chemical structure:

5 **Scaffold-NH-X(=V)-Functional entity precursor**

In which

X = -C-, -S-, -P-, -S(O)-, or -P(O)-, and

V = O, S, NH, or N-C₁-C₈ alkyl.

10

25. The method according to claim 24, wherein X is -C- and V is O.


Fig. 1




Fig. 2**Natural Base Pairs****Synthetic Base Pairs****Synthetic purine bases**

Fig. 3.

I = Inosine






Fig. 4




Fig. 5





Fig. 6



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 September 2003 (25.09.2003)

PCT

(10) International Publication Number
WO 03/078626 A3

(51) International Patent Classification⁷: **C07H 21/00**

(21) International Application Number: **PCT/DK03/00174**

(22) International Filing Date: **14 March 2003 (14.03.2003)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
PA 2002 0415 15 March 2002 (15.03.2002) DK
60/364,056 15 March 2002 (15.03.2002) US
02/00419 20 June 2002 (20.06.2002) DK
10/175,539 20 June 2002 (20.06.2002) US
60/434,427 19 December 2002 (19.12.2002) US

(71) Applicant (*for all designated States except US*): **NUEVO-LUTION A/S [DK/DK]**; Rønnegade 8, 5th floor, DK-2100 Copenhagen Ø (DK).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GOULIAEV, Alex, Haahr [DK/DK]**; Brøndsted 223, DK-3670 Veksø Sjælland (DK). **HO, Justin [US/DK]**; Mattæusgade 50, 3.-26, DK-1666 København V (DK). **FELDING, Jakob [DK/DK]**; Ordstruphøjvej 24, 1., DK-2920 Charlottenlund (DK). **SAMS, Christian [DK/DK]**; Jakob Dannefærdsvej 4A, 1., DK-1973 Frederiksberg C (DK). **PEDERSEN, Henrik [DK/DK]**; Frodesvej 24, DK-2880 Bagsværd (DK). **JENSEN, Kim, Birkebæk [DK/DK]**; Voldumvej 30C, DK-2610 Rødovre (DK). **HANSEN, Anders, Holm [DK/DK]**; Slotsfogedvej 3, st. th., DK-2400 København NV (DK). **LUNDORF, Mikkel, Dybro [DK/DK]**; Charlotte Munksvej 31, 2. tv., DK-2400 København NV (DK).

HUSEMOEN, Gitte, Nystrup [DK/DK]; Bregnerødgade 18, 1.th., DK-2200 København N (DK). **FRANCH, Thomas [DK/DK]**; Humlebækgade 14, st. tv., DK-2200 København N (DK). **THISTED, Thomas [DK/DK]**; Fjordskrænten 14, DK-3600 Frederikssund (DK).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
4 December 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **A BUILDING BLOCK CAPABLE OF TRANSFERRING A FUNCTIONAL ENTITY**

(57) Abstract: A building block having the dual capabilities of transferring the genetic information e.g. by recognising an encoding element and transferring a functional entity to a recipient reactive group is disclosed. The building block can be designed with an adjustable transferability taking into account the components of the building block. The building block may be used in the generation of a single complex or libraries of different complexes, wherein the complex comprises an encoded molecule linked to an encoding element. Libraries of complexes are useful in the quest for pharmaceutically active compounds.

WO 03/078626 A3

INTERNATIONAL SEARCH REPORT

PCT/DK 03/00174

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07H21/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 07734 A (HYBRIDON INC) 26 February 1998 (1998-02-26) figures	1-4, 8, 9, 12-20
X	US 6 326 478 B1 (CHERUVALLATH ZACHARIA S ET AL) 4 December 2001 (2001-12-04) column 17 -/-	1-4, 8, 9, 12-20

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

22 September 2003

Date of mailing of the international search report

06/10/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

de Nooy, A

INTERNATIONAL SEARCH REPORT

PCT/DK 03/00174

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WALDER J A ET AL: "COMPLEMENTARY CARRIER PEPTIDE SYNTHESIS: GENERAL STRATEGY AND IMPLICATIONS FOR PREBIOTIC ORIGIN OF PEPTIDE SYNTHESIS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, vol. 76, no. 1, January 1979 (1979-01), pages 51-55, XP000857351 ISSN: 0027-8424 the whole document	22
A	BRUICK R K ET AL: "TEMPLATE-DIRECTED LIGATION OF PEPTIDES TO OLIGONUCLEOTIDES" CHEMISTRY AND BIOLOGY, CURRENT BIOLOGY, LONDON, GB, vol. 3, no. 1, January 1996 (1996-01), pages 49-56, XP000856876 ISSN: 1074-5521 the whole document	22
A	US 5 693 773 A (AGRAWAL SUDHIR ET AL) 2 December 1997 (1997-12-02) figure 11	1
A	WO 00 14102 A (FUJISAWA KAZUHIKO ;JAPAN SCIENCE & TECH CORP (JP); NAKATANI KAZUHI) 16 March 2000 (2000-03-16) abstract	1

INTERNATIONAL SEARCH REPORT

PCT/DK 03/00174

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1-25 (in part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/DK 03/00174

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9807734	A	26-02-1998	AU	4156197 A	06-03-1998
			WO	9807734 A1	26-02-1998
US 6326478	B1	04-12-2001	AU	4977999 A	01-02-2000
			US	2002055623 A1	09-05-2002
			WO	0002896 A1	20-01-2000
			US	2003149260 A1	07-08-2003
			US	6399756 B1	04-06-2002
US 5693773	A	02-12-1997	AU	6255896 A	30-12-1996
			WO	9640710 A1	19-12-1996
WO 0014102	A	16-03-2000	JP	2000086692 A	28-03-2000
			WO	0014102 A1	16-03-2000

INTERNATIONAL SEARCH REPORT

International Application No. PCT/DK 03 00174

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-25 (in part)

Present claims 1-25 relate to an extremely large number of possible building blocks. In fact, the claims contain so many options that a lack of clarity within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. Moreover, support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found for only a very small proportion of the building blocks claimed. Consequently, the search has been carried out for those parts of the application which do appear to be clear and supported, namely those parts of the application relating to the building blocks of claim 1 where the complementing element is a nucleic acid or a derivative thereof as in claims 16 and 17 AND where the C-F connecting group is as defined in claim 15.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.